

Alternative Isoforms of TonEBP with Variable N-termini are Expressed in Mammalian Cells

Hyo Shin Kim, Sook Jin Son, Seon Nyo Kim, Yong Duk Kim, Kwang Jin Kim, Byeong Hwa Jeon, Jin Bong Park, and Sang Do Lee¹

Department of Physiology, College of Medicine, ¹Research Institute for Medical Sciences, Chungnam National University, Daejeon 301-131, Korea

Hypertonicity imposes a great deal of stress to cells since it causes rise in cellular ionic strength, which can be reduced by the accumulation of compatible osmolytes. TonEBP plays a central role in the cellular accumulation of compatible osmolytes via transcriptional stimulation of membrane transporters and aldose reductase. Alternatively spliced forms of TonEBP mRNA have previously been reported and two of them showed different transcriptional activity. In the present study, isoform-specific antibodies were produced to confirm the translation of the spliced mRNA to protein. TonEBP was immunoprecipitated by using anti-TonEBP antibody and then immunoblotted using anti-TonEBP or isoform specific antibodies to find out the expression profile of TonEBP isoforms in basal or stimulated condition. From these results, we conclude that all TonEBP isoforms are expressed in mammalian cells and their expression patterns are not same in every cells.

Key Words: TonEBP, Hypertonicity, Alternative splicing

INTRODUCTION

TonEBP plays a central role in the cellular accumulation of compatible osmolytes via transcriptional stimulation of membrane transporters and aldose reductase (Ko et al, 1997; Rim et al, 1998; Miyakawa et al, 1999a; Cha et al, 2001). The critical importance of TonEBP in mediating adaptation to hyperosmotic stress within the kidney has been demonstrated by targeted disruption of the TonEBP gene in mouse resulting in a null allele (Lopez-Rodriguez et al, 2004). Although the majority of homozygous TonEBP^{-/-} animals die late in gestation or perinatally, approximately 3% of the homozygous progeny survive past weaning. These animals exhibit marked atrophy of the renal medulla that is associated with impaired activation of several TonEBP-regulated osmoprotective genes. Consistent with this result, transgenic over-expression of a dominant inhibitory form of TonEBP in the epithelial cells of the renal collecting tubules results in an impaired capacity to concentrate urine, leading to progressive hydronephrosis, that is also associated with defects in the expression of osmo-regulatory genes (Lam et al, 2004). Therefore, as expected from the unique physiology of mammalian kidney, which is characterized by the generation of a hyperosmotic environment within the renal medulla during antidiuresis, the osmotic stress response pathway which is defined by the TonEBP

transcription factor is clearly essential for normal renal function.

TonEBP plays also a role in T-cell growth, cell migration and invasion. TonEBP is activated by T-cell receptor activation (Trama et al, 2000), and murine transgenic line expressing a dominant negative form of TonEBP results in lymphoid hypocellularity and impaired T lymphocyte growth (Trama et al, 2002). TonEBP is also activated by the cross-linking of $\alpha 6 \beta 4$ integrin, and the expression of an inhibitory form of TonEBP in cancer cells impairs cell invasion (Jauliac et al, 2002). In addition, overexpression of TonEBP results in enhanced cell migration.

The human TonEBP gene is made up of 16 exons over 130 kbp (Maouyo et al, 2002). Due to alternative splicing involving exons 2~4, three different polypeptides are predicted. All the deduced splice isoforms share the same 1,455 amino acids of the "a" form (Miyakawa et al, 1999b) that includes the DNA binding domain. Other isoforms have additional 76~96 amino acids at the N-terminus. In the present study, we produced isoform-specific antibodies and showed that the isoforms are expressed as proteins.

METHODS

DNA constructs

All constructs were generated using standard cloning procedures and verified by restriction enzyme analysis and

ABBREVIATIONS: TonEBP, Tonicity-responsive enhancer (TonE) binding protein; RHD, Rel Homology domain.

Corresponding to: Sang Do Lee, Department of Physiology, College of Medicine, Research Institute for Medical Sciences, Chungnam National University, 6, Munhwa-dong, Jung-gu, Daejeon 301-131, Korea. (Tel) 82-42-580-8220, (Fax) 82-42-585-8440, (E-mail) sdlee@cnu.ac.kr


DNA sequencing. Each full-length human TonEBP cDNA was cloned into a mammalian expression vector pCMV-Tag2 (Stratagene, La Jolla, CA) which allows the expression of TonEBP fused with FLAG epitope. Each Y1 fragments of TonEBP were generated by PCR or using restriction enzyme and cloned into pCMV-Tag2.

Cell culture and transfection

COS7 cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS; Gemini Bio-Products, Woodland, CA), 100 μ g/ml penicillin, and 100 μ g/ml streptomycin (Life Technologies Inc. Grand Island, NY). HeLa cells were maintained in DMEM supplemented with 10% horse serum (Quality Biological, Inc. Gaithersburg, MD), 100 μ g/ml penicillin, and 100 μ g/ml streptomycin. Jurkat cells were maintained in RPMI 1640 supplemented with 10% FBS, 100 μ g/ml penicillin, and 100 μ g/ml streptomycin. mIMCD cells were maintained in DMEM/F-12 supplemented with 10% FBS, 100 μ g/ml penicillin, and 100 μ g/ml streptomycin. CHO cells were maintained in MEM-alpha supplemented with 10% FBS, 100 μ g/ml penicillin, and 100 μ g/ml streptomycin. Cells were transfected using Lipofectamine 2000 as instructed by the manufacturer (Life Technologies Inc. Grand Island, NY).

Antibodies, Immunoprecipitation, and Immunoblot

Each peptides corresponding to exon 2 (for anti-E2 antibody) or N-terminus of exon 3 (for anti-E3 antibody) were synthesized, purified, and then used to raise antibodies in rabbits through a commercial service (Covance, Denver, PA). For immunoprecipitation, cells were washed once with ice-cold phosphate-buffered saline (PBS) and lysed in lysis buffer containing 50 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 1 mM dithiothreitol, and protease inhibitor cocktail (Roche Molecular Biochemicals, Indianapolis, IN) for 30 min at 4°C. Lysates were centrifuged for 5 min at 15,000 g to pellet insoluble materials. Resulting supernatants (whole cell extracts) were incubated for 60 min with 10 μ l of anti-TonEBP serum and 40 μ l of protein A agarose (Life Technologies Inc. Grand Island, NY) to immunoprecipitate endogenous TonEBP at 4°C. The immune complexes were then washed four times in lysis buffer, resuspended in the sample buffer [62.5 mM Tris (pH 6.8), 25% glycerol, 2% SDS, 0.01% bromophenol blue, 5% 2-mercaptoethanol], and heated at 95°C for 5 min. For immunoblot, whole cell extracts or immune complexes were resolved on SDS-polyacrylamide gels and transferred to nitrocellulose membrane. Nonspecific binding sites were blocked with 5% nonfat milk in TTBS [25 mM Tris (pH 7.4), 137 mM NaCl, 3 mM KCl, 0.05% Tween20] for 30 min at room temperature. The membrane was incubated with 1 : 200 dilution of anti-E3 antibody, with 1 : 600 dilution of anti-E2 antibody, with 1 : 2,000 dilution of anti-FLAG M5 antibody (Sigma, Saint Louis, MO) or anti-TonEBP antibody in blocking solution for 60 min at room temperature. The membrane was then incubated with 1 : 5,000 dilution of anti-mouse IgG or anti-rabbit IgG conjugated with alkaline phosphatase in the same way as described above (Jackson Immunoresearch, West Grove, PA). Alkaline phosphatase activity was visualized using a commercial kit (Sigma, Saint Louis, MO).



▲ amino acid sequences of exon 2

SLKLHPSQTFHRAGLLE

Exon2-specific => α -E2

▲ amino acid sequences of exon 3

SVYDLLPKELQLPPPRETSVASMSTSGGEAGSPPPAVVAA

Exon3-specific => α -E3

Fig. 1. Schematic representation of TonEBP isoform, based on previous reports. Amino acid sequences corresponding to exon 2 and exon 3 of TonEBP are shown. Each peptides used in antibody production is underlined.

RESULTS

We previously reported that widespread alternative splicing in exons 2~4 was detected during development as well as in different adult tissues. Therefore, it is highly possible that different polypeptides with different lengths at the N-terminus might be produced. Two of the isoforms have been shown to be different in their ability to stimulate transcription (Maouyo et al, 2002). In the present study, therefore, isoform-specific antibodies were prepared to confirm the translation of the spliced mRNA to protein. Each peptides corresponding to exon 2 for TonEBP-d isoform or N-terminus of exon 3 for TonEBP-c and -d isoform were synthesized, and they were then used to raise antibodies as described in materials and method (Fig. 1). Affinity of each antibodies produced was tested using cell lysates which express FLAG-fused N-terminal fragment of TonEBP-d isoform (FLAG-Yd1) as antigen (data not shown). FLAG-fused full-length or Y1 fragments of each TonEBP isoforms were expressed in COS7 cells and then used for characterization of each antibodies (Fig. 2). Anti-FLAG antibody or anti-TonEBP antibody which recognizes all isoforms of TonEBP were used as a positive control to confirm the expression of each protein. TonEBP-c isoform (FLAG-Yc1 or FLAG-TonEBP-c) was detected by anti-E3 antibody, but not by anti-E2 antibody, whereas TonEBP-d isoform (FLAG-Yd1 or FLAG-TonEBP-d) was detected by both antibodies as expected. From these results, isoform specificity of each antibodies was confirmed; that is to say, anti-E2 antibody detects only TonEBP-d isoform, whereas anti-E3 antibody detects both TonEBP-c and -d isoforms.

To elucidate the expression profile of TonEBP isoforms in basal or stimulated condition, endogenous TonEBP was immunoprecipitated by using anti-TonEBP antibody and then immunoblotted using anti-TonEBP antibody or isoform specific antibodies (Fig. 3). In Jurkat T-cells and HeLa cells, TonEBP was not detected by both anti-E2 antibody and anti-E3 antibody, indicating that only TonEBP-a iso-




Fig. 2. Antibodies are specific to each TonEBP isoform. FLAG-fused full-length (B) or Y1 fragments (A) of each TonEBP isoform were used for characterization of each antibody. Anti-FLAG antibody and anti-TonEBP antibody were used as a positive control to confirm the expression of each protein.

form was expressed. On the other hand, TonEBP in mIMCD cells was detected by all antibodies used, suggesting that TonEBP-d isoform was expressed. TonEBP in CHO cells was detected by anti-E3 antibody, but not by anti-E2 antibody, indicating that TonEBP-c isoform was expressed. From these results, we can conclude that all TonEBP isoforms are expressed in mammalian cells and their expression patterns are variable, depending on cell types.

DISCUSSION

The amino acid sequence and gene structure of TonEBP are highly homologous between humans and mice. A genome database search reveals that Rel-like transcription factors are present in *Drosophila* and mammals, but not in *Caenorhabditis elegans*, yeast, and plants (Riechmann et al, 2000). *Drosophila* has one homolog of TonEBP, named as MESR1 (GenBank accession no. AF195496). MESR1 was identified as a modifier of RAS1 signaling involved in eye development (Huang & Rubin, 2000).

We and others previously detected widespread alternative splicing in exon 2~4 during mouse development and in several cell lines (Maouyo et al, 2002; Kim et al, 2005). Consequently, four different mRNA and three possible polypeptides with different lengths at the N-terminus are expected to be produced (Fig. 1). TonEBP-c and TonEBP-d isoform have an additional 76 and 94 amino acids, respectively, at the N-terminus of the TonEBP-a, which is




Fig. 3. Each isoforms of TonEBP are expressed in several mammalian cells. Each cell line was treated with 100 mM NaCl for 16 hours. TonEBP was immunoprecipitated from whole cell lysates using anti-TonEBP antibody. Immune complexes were then resolved on 6% SDS-polyacrylamide gels and probed using each antibody. I or H represents isotonic or hypertonic conditions, respectively.

the shortest. Compared with adult human brain, the abundance of the TonEBP-a transcript is much higher in the human fetal brain than other spliced isoforms (Dalski et al, 2000). In other tissue, there is little difference in the relative abundance of TonEBP isoforms between fetuses and adults. By using isoform-specific antibodies, we found that the expression patterns of TonEBP isoforms were different in several cell lines (Fig. 3).

Alternative splicing generates multiple protein isoforms from a single gene and contributes to the diversity of the eukaryotic proteome (Black, 2000; Graveley, 2001; Caceres & Kornblihtt, 2002). By varying the inclusion of functionally significant sequences, such as CoRRN boxes and their flanking sequences, it is possible to generate corepressor isoforms that differ in their affinity for transcription factors (Goodson et al, 2005). This is likely to contribute to the refinement of the regulation of gene expression. Insertion of N-terminal 76 and 94 amino acids in the TonEBP-c and TonEBP-d isoform results in different basal and hypertonicity-induced TonEBP activity (Maouyo et al, 2002; Kim et al, 2005), and TonEBP-c shows the highest activity in both isotonic and hypertonic conditions.

Another insertion of 18 amino acids to the TonEBP-c produces TonEBP-d, which reduces TonEBP activity, although TonEBP-d activity is still higher than TonEBP-a. Expression of TonEBP is known to be induced by T-cell receptor and integrin signaling as well as hypertonicity. Phorbol-ester-induced macrophagic differentiation in U937 cells leads to surface expression of Mac-1 integrin and its activation as well (Hamada & Utiyama, 2005). Expression of TonEBP is also induced during PMA-induced macrophagic differentiation, although it is not known whether Mac-1 integrin is involved in this pathway (data not shown). These observations suggest that different expression profiles of TonEBP isoforms in different tissues and in response to different stimulations, such as hypertonicity, T-cell receptor, and integrin, may be important for the regulation of TonEBP activity.

ACKNOWLEDGMENT

This work was supported in part by research fund of Chungnam National University in 2004 and in part by the Korea Research Foundation Grant Funded by the Korean Government (R08-2003-000-10599-0).

REFERENCES

- Black DL. Protein diversity from alternative splicing: a challenge for bioinformatics and post-genome biology. *Cell* 103: 367–370, 2000
- Caceres JF, Kornblihtt AR. Alternative splicing: multiple control mechanisms and involvement in human disease. *Trends Genet* 18: 186–193, 2002
- Cha JH, Woo SK, Han KH, Kim YH, Handler JS, Kim J, Kwon HM. Hydration status affects nuclear distribution of transcription factor TonEBP in rat kidney. *J Am Soc Nephrol* 12: 2221–2230, 2001
- Dalski A, Wagner HJ, Schwinger E, Zuhlke C. Quantitative PCR analysis of different splice forms of NFAT5 revealed specific gene expression in fetal and adult brain. *Mol Brain Res* 83: 125–127, 2000
- Goodson ML, Jonas BA, Privalsky ML. Alternative mRNA splicing of SMRT creates functional diversity by generating corepressor isoforms with different affinities for different nuclear receptors. *J Biol Chem* 280: 7493–7503, 2005
- Graveley BR. Alternative splicing: increasing diversity in the proteomic world. *Trends Genet* 17: 100–107, 2001
- Hamada K, Utiyama H. Functional cytoplasmic domains of the Mac-1 integrin receptor in phorbol ester-treated U937 cells. *Biochem Biophys Res Commun* 335: 858–864, 2005
- Huang AM, Rubin GM. A misexpression screen identifies genes that can modulate RAS1 pathway signaling in *Drosophila melanogaster*. *Genetics* 156: 1219–1230, 2000
- Jauliac S, Lopez-Rodriguez C, Shaw LM, Brown LF, Rao A, Toker A. The role of NFAT transcription factors in integrin-mediated carcinoma invasion. *Nat Cell Biol* 4: 540–544, 2002
- Kim HS, Park JI, Lee SD. Expression and functional differences of the alternative spliced TonEBP isoforms. *Korean J Gerontol* 15: 17–25, 2005
- Ko BCB, Ruepp B, Bohren KM, Gabbay KH, Chung SSM. Identification and characterization of multiple osmotic response sequences in the human aldose reductase gene. *J Biol Chem* 272: 16431–16437, 1997
- Lam AK, Ko BC, Tam S, Morris R, Yang JY, Chung SK, Chung SS. Osmotic response element binding protein (OREBP) is an essential regulator of urine concentrating mechanism. *J Biol Chem* 279: 48048–48054, 2004
- Lopez-Rodriguez C, Antos CL, Shelton JM, Richardson JA, Lin F, Novobrantseva TI, Bronson RT, Igarashi P, Rao A, Olson EN. Loss of NFAT5 results in renal atrophy and lack of tonicity-responsive gene expression. *Proc Natl Acad Sci USA* 101: 2392–2397, 2004
- Maouyo D, Kim JY, Lee SD, Wu Y, Woo SK, Kwon HM. Mouse TonEBP-NFAT5: expression in early development and alternative splicing. *Am J Physiol* 282: F802–F809, 2002
- Miyakawa H, Rim JS, Handler JS, Kwon HM. Identification of the second tonicity-responsive enhancer for the betaine transporter (BGT1) gene. *Biochim Biophys Acta* 1446: 359–364, 1999a
- Miyakawa H, Woo SK, Dahl SC, Handler JS, Kwon HM. Tonicity responsive enhancer binding protein, a rel-like protein that stimulates transcription in response to hypertonicity. *Proc Natl Acad Sci USA* 96: 2538–2542, 1999b
- Riechmann JL, Heard J, Martin G, Reuber L, Jiang CZ, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, Creelman R, Pilgrm M, Broun P, Zhang JZ, Ghandehari D, Sherman BK, Yu GL. Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290: 2105–2110, 2000
- Rim JS, Atta MG, Dahl SC, Berry GT, Handler JS, Kwon HM. Transcription of the sodium/myo-inositol cotransporter gene is regulated by multiple tonicity-responsive enhancers spread over 50 kilobase pairs in the 5-flanking region. *J Biol Chem* 273: 20615–20621, 1998
- Trama J, Go WY, Ho SN. The osmoprotective function of the NFAT5 transcription factor in T cell development and activation. *J Immunol* 169: 5477–5488, 2002
- Trama J, Lu Q, Hawley RG, Ho SN. The NFAT-related protein NFATL1 (TonEBP/NFAT5) is induced upon T cell activation in a calcineurin-dependent manner. *J Immunol* 165: 4884–4894, 2000