

RESEARCH COMMUNICATION

HIF-1 α siRNA and Cisplatin in Combination Suppress Tumor Growth in a Nude Mice Model of Esophageal Squamous Cell Carcinoma

Hong-Ying Liao^{1*}, Gui-Ping Wang², Li-Jia Gu¹, Shao-hong Huang¹, Xiu-Ling Chen¹, Yun Li¹, Song-Wang Cai¹

Abstract

Introduction: The esophagus squamous cell carcinoma (ESCC) is one of the most deadly malignances, and a current challenge is the development of effective therapeutic agents. Our present work addressed the effect of HIF-1 α siRNA alone or in combination with cisplatin on the growth of ESCC in nude mice. **Materials and Methods:** Xenografts were established by inoculating ESCC TE-1 cells in nude mice, and transplanted tumors were treated with HIF-1 α siRNA, cisplatin alone or together. Growth was assessed by measuring tumor volume. HIF-1 α mRNA and protein expression were detected using RT-PCR and immunohistochemistry, respectively. Apoptosis of ESCC TE-1 cells was analyzed by flow cytometry. **Results:** In our nude mice model, HIF-1 α siRNA effectively inhibited the growth of transplanted ESCC, downregulating HIF-1 α mRNA and protein expression, and inducing ESCC TE-1 cell apoptosis. Notably when combined with cisplatin, HIF-1 α siRNA showed synergistic interaction in suppressing tumor growth. Furthermore, the proportion of apoptotic cells in HIF-1 α siRNA plus cisplatin group was significantly higher than that in cisplatin or HIF-1 α siRNA-treated groups ($P < 0.05$). **Conclusions:** Down-regulated HIF-1 α expression induced by siRNA could effectively suppress the growth of transplanted ESCC *in vivo*. HIF-1 α siRNA could enhance the cytotoxicity of cisplatin, which suggests that a combination of these two agents may have potential for therapy of advanced ESCC.

Keywords: RNA interference - HIF-1 α - cisplatin - esophageal squamous cell carcinoma

Asian Pacific J Cancer Prev, 13, 473-477

Introduction

Squamous cell carcinoma is the main histological subtype of esophageal cancers in China. Clinically, cisplatin-based chemotherapy is the main approach for many patients with advanced esophageal cancer. Most patients with advanced esophageal cancer experienced chemotherapy failure, however, mechanisms for chemotherapeutic resistance have not been completely understood (Bazarbashi et al., 2008; Cho et al., 2008; Takashima et al., 2008). The transcription factor hypoxia inducible factor-1 (HIF-1), one heterodimer composed of HIF-1 α and HIF-1 β , is a critical mediator of the response to hypoxia. Recent studies revealed that hypoxia was associated with poor responses to chemotherapy in esophageal cancers (Koukourakis et al., 2001). More recently, evidences showed that increased expression of HIF-1 α was closely correlated with the growth, invasion and resistance to chemotherapeutic agents of human cancer cells including ESCC. In contrast, the loss of HIF-

1 α activity or suppression of HIF-1 α overexpression can effectively inhibit tumor growth, suggesting that HIF-1 α is a potent therapeutical target for ESCC (Matsuyama et al., 2005; Tzao et al., 2008).

RNA interference (RNAi), which is a sequence-specific gene-silencing method induced by double-stranded RNA (dsRNA), can precisely regulate or silence the expression of target genes. Numerous studies in animal models of human disease demonstrate the potential application of RNAi therapeutics (Sithanandam et al., 2005; Yonesaka et al., 2006; Angaji et al., 2010). Previous studies have shown that siRNA targeting HIF-1 α can induce apoptosis of some human tumor cells and suppress the tumor growth (Chen and Yu, 2009). More recently, a study reported that HIF-1 α siRNA could inhibit the proliferation of ESCC TE-1 cells *in vitro* (Wu et al., 2007). However, little is known about the effect of HIF-1 α siRNA alone or combination with cytotoxic agents on ESCC *in vivo*. In this study, we demonstrated that down-regulation of HIF-1 α expression using HIF-1 α siRNA alone or in combination with

¹Department of Thoracic Surgery, Clinical Research Center of Chest Tumor, the Third Affiliated Hospital, Sun Yat-sen University,

²Department of Pharmacy, Health College, Guangzhou Medical University, Guangzhou, China *For correspondence: Hylmed1996@hotmail.com

cisplatin induced ESCC TE-1 cell apoptosis and inhibited the growth of transplanted ESCC in nude mice. Our results suggest that HIF-1 α -targeting therapy may be a promising strategy to increase the efficacy of chemotherapy drugs for the treatment of esophagus cancers.

Materials and Methods

Cell culture and establishment of animal model

The BALB/c nude mice (28 \pm 3d, male and female) were purchased from the Experimental Animal Center of Sun Yat-sen University, and maintained under specific pathogen-free conditions. Human ESCC cell line TE-1 was provided by Shanghai Cell Bank of Chinese Academy of Science, and cultured in RPMI 1640 (Gibco, USA) containing 15% calf serum, 100 U/ml penicillin and 100 mg/L streptomycin at 37°C in a humidified atmosphere containing 5% CO₂. About 1 \times 10⁷ TE-1 cells were subcutaneously injected into the right upper limb of nude mice. When the tumor size was approximately 0.8 cm in diameter, the inhibition experiments were carried out. All animal experiments were approved by the animal care committee of SUN Yat-sen University and were performed using the criteria for humane care as outlined in the "Guide for the Care and Use of Laboratory Animals" published by the National Institutes of Health.

Treatment

siRNA targeting HIF-1 α was designed and validated by Boya Bio-engineering Company (Boya, China). The antisense sequences of siRNA targeting HIF-1 α was 5'-AGTTCACCTGAGCCTAATA-3'. Twenty nude mice were randomly assigned into four groups, and treated with saline, cisplatin (20 μ g/ml), HIF-1 α siRNA (20 μ mol/L) and HIF-1 α siRNA (20 μ g/ml) plus cisplatin (20 μ g/ml), respectively. The saline or cisplatin was injected into the center of the tumors, and HIF-1 α siRNA was transfected into nude mice with the lipofection according to the manufacturer's instructions of Lipofectamine 2000 (Invitrogen, USA). The longest (a) and shortest (b) diameters were measured every week and the volume of tumor was calculated as the formula $V=a^2 \times b/2$. Three weeks later, the nude mice were sacrificed, and tumor tissues were removed and weighted. Tumor growth inhibition rate (IR) was calculated according to the following formula: $IR(\%) = (1 - \text{mean weight experimental group} / \text{mean weight control group}) \times 100\%$ (Kim et al., 2011).

Determination of HIF-1 α mRNA with RT-PCR

The total RNA in tumor tissues was extracted according to the manufacturer's protocol and stored in -80°C for use. Subsequently, one-step RT-PCR was performed to detect HIF-1 α mRNA according to manufacturer's instructions (Wang et al., 2011). The primers for HIF-1 α and GAPDH (as internal normalization control) were as follows: HIF-1 α : 5'-CAGCAACCAGGTGACCGTG-3' (forward) and 5'-TGCTGCCTTGTATGGGAGCATT-3' (reverse); GAPDH: 5'-CGGAGTCAACGGATTTGGTGGTAT-3' (forward) and 5'-AGCCTTCTCCATGGTGGTGAAGAC-3' (reverse). The reaction mixture contained 3 μ l of

template RNA, 2 μ l of each primer, 17.8 μ l of DEPC treated water, 25 μ l of 2 \times buffer and 0.2 μ l of RT-Taq polymerase (5U/ μ l). Reverse transcription from RNA to cDNA was performed at 50°C for 50min. The DNA was denatured at 94°C for 2 min, following by 30 cycles of amplification. Each cycle consisted of denaturation at 94°C for 45s, annealing at 55°C for 1min, and extension at 72°C for 1min, with an additional extension at 72°C for 7min. The products were stored at 4°C. A total of 5 μ l of PCR products was added into 1 μ l 6 \times loading buffer which was loaded onto 1.5% agarose gel for electrophoresis. The RNA bands were visualized under UV light and the optical density was analyzed which was normalized by the density of GAPDH.

Detection of HIF-1 α protein with immunohistochemistry

The immunohistochemistry was performed according to the protocol of S-P kit (Invitrogen, USA). The primary antibody was mouse anti-human polyclonal antibody (Santa, USA), and the secondary antibody was included in the kit. The slides were observed under microscope. Light brown in the cytoplasm suggested weakly positive, brown was regarded as strong positive, and primary color indicated negative (Matsuyama et al., 2005; Tzao et al., 2008). The optical density was analyzed with automatic image analysis system (Kontron IBAS 2.5, Germany) and positive unit (PU) was obtained.

Analysis of apoptosis with flow cytometry

Cell apoptosis assays were done as previously described (Chen et al., 2009; Wang et al., 2011). In brief, the tumor tissues were treated with 0.25% trypsin and 0.02% EDTA. Cell suspension was collected and centrifuged at 1,000 rpm for 5 min. Supernatant was removed and cells were rinsed with PBS thrice. Then, these cells were suspended in 100 μ l binding buffer at a final of 1 \times 10⁶/ml, and incubated with 10 μ l Annexin V-FITC (Invitrogen, USA) for 30 min. Subsequently, 5 μ l PI was added and incubated away from light at room temperature for five minutes. After that, the cells were directly analyzed by FACScan and evaluated by the CellQuest program (Becton Dickinson, USA).

Statistical analysis

The data were presented as mean \pm SD, and analyzed using SPSS 15.0 software package. One-way analysis of variance (ANOVA) followed by a post hoc LSD-t or Dunnett's test was performed to determine the significance of the differences in multiple comparisons. A value of $P < 0.05$ was considered statistically significant.

Results

Combination of HIF-1 α siRNA with cisplatin significantly suppressed the growth of transplanted ESCC

To test the ability of HIF-1 α siRNA alone or in combination with cisplatin to suppress the growth of ESCC, 1 \times 10⁷ TE-1 cells were subcutaneously injected into the right upper limb of nude mice. Two weeks later, the tumor size was approximately 0.6~0.8 cm in diameter. No significant difference in tumor volume was found




Figure 1. Antitumor Effect of HIF-1 α siRNA, Cisplatin Alone or Together Against ESCC in Nude Mice. TE-1 cells (1×10^7) were subcutaneously injected into the right upper limb of BALB/c nude mice. Two weeks later, nude mice were treated with indicated agents. Tumor volume were measured to assess the antitumor effect of indicated agents weekly. (A) HIF-1 α siRNA, or HIF-1 α siRNA in combination with cisplatin strongly inhibited the growth of the tumor (B,C). Each bar represents the means \pm SD (n = 3), *P < 0.05




Figure 2. Effect of HIF-1 α siRNA or in Combination with Cisplatin on HIF-1 α mRNA Expression in ESCC Tissue. HIF-1 α mRNA was detected by RT-PCR. Relative mRNA expression of HIF-1 α was calculated to evaluate the ability of indicated agents to suppress HIF-1 α mRNA expression (A). When compared with control group, HIF-1 α mRNA expression in HIF-1 α siRNA alone or together with cisplatin group was significantly suppressed (*P < 0.05). Each bar represents the mean \pm SD (n=3). PCR products was analyzed and identified with agarose gel electrophoresis (B). m: Marker; a: Saline; b: Cisplatin; c: HIF-1 α siRNA; d: Cisplatin+HIF-1 α siRNA.

between groups before treatment ($p > 0.05$). After nude mice xenograft were treated with cisplatin, HIF-1 α siRNA alone or together, the growth of tumor was inhibited to different extents. As showed in Figure 1, a statistically significant reduction in the speed of tumor growth was observed in nude mice treated with HIF-1 α siRNA alone or in combination with cisplatin compared with that in control group (Figure 1A,C). When compared with HIF-1 α siRNA or cisplatin alone, the combination of HIF-1 α siRNA and cisplatin strongly suppressed the growth of transplanted ESCC, the inhibition rate was upto 52.4% which was significantly higher than that of cisplatin- or HIF-1 α siRNA-treated group ($P < 0.05$) (Figure 1B).

HIF-1 α siRNA downregulated HIF-1 α mRNA expression

To assess the potency of the HIF-1 α siRNA to block HIF-1 α expression, mRNA expression level of HIF-1 α in transplanted ESCC tissues was analysed by RT-PCR. As shown in Figure 2, when compared to saline or




Figure 3. HIF-1 α Protein Expression in Tumor Tissue from Transplanted ESCC. HIF-1 α protein Expression was determined by immunohistochemistry. The HIF-1 α was stained in brown color in the photographs taken at the magnifications of 400 \times (A) Immunohistochemical staining revealed that HIF-1 α siRNA alone or together with cisplatin significantly inhibited HIF-1 α expression. (B) Each bar represents the mean \pm SD (n = 3), *P < 0.05




Figure 4. Effect of HIF-1 α siRNA or in Combination with Cisplatin on Cell Apoptosis Quantitated Using Annexin-V/PI Detection. The tumor tissues were obtained from ESCC treated by HIF-1 α siRNA, cisplatin alone or together in nude mice. ESCC TE-1 cells were harvested and analyzed by flow cytometry. (A) The combination of HIF-1 α siRNA with cisplatin significantly induced the apoptosis of TE1 cells. (B) Each bar represents the means \pm SD (n = 3), *P < 0.05

cisplatin treated nude mice, HIF-1 α siRNA alone or in combination with cisplatin significantly down-regulated the expression of HIF-1 α mRNA ($P < 0.05$). However, no difference in mRNA expression of HIF-1 α was observed between HIF-1 α siRNA and HIF-1 α siRNA combined with cisplatin group ($P > 0.05$). These results indicated that HIF-1 α siRNA could effectively inhibit HIF-1 α mRNA expression.

HIF-1 α siRNA suppressed HIF-1 α protein expression

To investigate the effect of HIF-1 α siRNA on the HIF-1 α protein expression, the immunohistochemistry analysis was performed. The results were shown in Fig. 3. HIF-1 α protein expression was not suppressed in cisplatin group in comparison to saline control ($P > 0.05$). However, HIF-1 α protein expression was significantly inhibited in ESCC tissues treated with HIF-1 α siRNA or HIF-1 α siRNA plus cisplatin when compared with that in saline or cisplatin treated group ($P < 0.05$). But, the reduction of HIF-1 α protein expression showed no difference between HIF-1 α siRNA and HIF-1 α siRNA plus cisplatin group

($P > 0.05$).

HIF-1 α siRNA induced ESCC TE-1 cell apoptosis

Flow cytometric experiments were performed to determine if the growth inhibition of transplanted ESCC treated with HIF-1 α siRNA alone or in combination with cisplatin was related with the apoptosis of ESCC TE-1 cell. Annexin-V and PI staining were used to detect apoptotic cells including early-stage apoptosis (FITC+/PI-) and late-stage apoptosis or necrotic cells (FITC+/PI+) (Wang et al., 2011). As shown in Figure 4, 28%~34% of apoptotic cells were observed in cisplatin or HIF-1 α siRNA-treated group as compared to about 6% apoptotic cells in saline control ($P < 0.05$). When HIF-1 α siRNA combined with cisplatin, the percentage of apoptotic cells (upto 60%) was significantly higher than that in cisplatin or HIF-1 α siRNA-treated group ($P < 0.05$).

Discussion

In most human cancers including ESCC, over-expression of HIF-1 α promotes tumor growth, angiogenesis and disease progression. Thus, targeting HIF-1 α may serve as a viable anticancer strategy to be used in clinics (Kurokawa et al., 2003; Kimura et al., 2004; Matsuyama et al., 2005). Herein, using the strategy of silencing HIF-1 α with targeted siRNA, we identified that HIF-1 α siRNA could effectively inhibit the growth of transplanted ESCC in nude mice, down-regulate the HIF-1 α mRNA and protein expression, and induce ESCC TE-1 cell apoptosis. Interestingly, when combined with the commonly-prescribed cisplatin, HIF-1 α siRNA showed synergistic interaction in inhibiting the tumor growth. These results suggested that combination of siRNA targeting HIF-1 α and existing chemotherapeutic agents maybe a promising approach in the treatment of advanced or chemotherapy-resistant ESCC.

A deficiency in apoptosis is a well-known hallmark of many cancers. Many studies validated that HIF-1 α is concerned with apoptosis, however HIF-1 α possesses dual functions in mediating apoptosis of tumour cells, i.e, anti-apoptotic and pro-apoptotic function (Flamant et al., 2010; Khan et al., 2011). The overall effect of HIF-1 α on cell apoptosis is still not fully understood, maybe depend on microenvironment in tumor, especially hypoxic condition. HIF-1 α may induce cell apoptosis through enhancing the stabilization of p53 protein under severe hypoxic condition (Hammond and Giaccia, 2005). Pancreatic cancer cell lines with constitutively expression of HIF-1 α showed a protective role in limiting hypoxia-induced apoptosis. As a fast-growing cancer, ESCC has a hypoxic microenvironment and metabolic disturbance. Hypoxia suppresses the apoptosis of cancer cells, resulting in the resistance to chemotherapeutic agents and even treatment failure (Matsuyama et al., 2005; Tzao et al., 2008). As shown in our results (Figure 4), HIF-1 α siRNA alone or in combination with cisplatin strongly induced apoptosis of ESCC TE-1 cell, which resulted in the growth suppression of transplanted ESCC in nude mice.

Several small molecular inhibitors (e.g. temsirolimus,

everolimus, bevacizumab, rencarex, cisplatin, etc) have been identified to decrease the level of HIF-1 α mRNA and protein (Rapisarda et al., 2004; Duyndam et al., 2007; Seeber et al., 2010). Moreover, some of these HIF-1 inhibitors are subject of clinical trials at present. Specific siRNA has been showed to effectively reduce HIF-1 α expression (Welsh et al., 2004). In the present study, our data revealed that HIF-1 α siRNA we designed could effectively suppress the HIF-1 α mRNA and protein expression in vivo (Figure 2 and 3). Suppression of tumor growth could be directly caused by the down-regulation of HIF-1 α mRNA and protein in ESCC tissues. However, no significant difference in the expression of HIF-1 α mRNA and protein was observed between HIF-1 α siRNA and HIF-1 α siRNA plus cisplatin group, which might be resulted from the effect of two agents combination masked by the cytotoxicity of cisplatin.

In conclusion, our study demonstrated that inhibition of HIF-1 α is a potential strategy for the treatment of ESCC. Since HIF-1 α siRNA enhanced the response of cancer cells to cisplatin, a combined therapy of HIF-1 α siRNA and cisplatin or other chemotherapeutic agents maybe a promising approach for ESCC. However, the specific delivery, the stability and transfection efficiency of siRNA into target tissues remains a challenge (Sithanandam et al., 2005; Seeber et al., 2010). The anti-tumor activity HIF-1 α siRNA or in combination with cisplatin in ESCC need to be verified in subsequent trials, subjecting to more in-depth studies.

Acknowledgements

This work was supported by the Natural Science Foundation for Doctoral Initiating Research of Guangdong Province (No. 06300768). We greatly appreciate Dr. Qianglin Duan from Tongji University, for manuscript revision.

References

- Angaji SA, Hedayati SS, Poor RH, et al (2010). Application of RNA interference in treating human diseases. *J Genet*, **89**, 527-37.
- Bazarbashi S, Abdelsalam M, Amin T, et al (2008). Phase II trial of neoadjuvant cisplatin, 5-fluorouracil and interferon-alpha in operable squamous cell carcinoma of the esophagus. *Chemotherapy*, **54**, 315-22.
- Chen C, Yu Z (2009). siRNA targeting HIF-1 α induces apoptosis of pancreatic cancer cells through NF-kappaB-independent and -dependent pathways under hypoxic conditions. *Anticancer Res*, **29**, 1367-72.
- Chen Y, Lu Y, Lu C, et al (2009). Beclin-1 expression is a predictor of clinical outcome in patients with esophageal squamous cell carcinoma and correlated to hypoxia-inducible factor (HIF)-1 α expression. *Pathol Oncol Res*, **15**, 487-93.
- Cho SH, Shim HJ, Lee SR, et al (2008). Concurrent chemoradiotherapy with S-1 and cisplatin in advanced esophageal cancer. *Dis Esophagus*, **21**, 697-703.
- Duyndam MC, van Berkel MP, Dorsman JC, et al (2007). Cisplatin and doxorubicin repress Vascular Endothelial Growth Factor expression and differentially down-regulate

- Hypoxia-inducible Factor I activity in human ovarian cancer cells. *Biochem Pharmacol*, **74**, 191-201.
- Flamant L, Notte A, Ninane N, et al (2010). Anti-apoptotic role of HIF-1 and AP-1 in paclitaxel exposed breast cancer cells under hypoxia. *Mol Cancer*, **9**, 191.
- Hammond EM, Giaccia AJ (2005). The role of p53 in hypoxia-induced apoptosis. *Biochem Biophys Res Commun*, **331**, 718-25.
- Khan MN, Bhattacharyya T, Andrikopoulos P, et al (2011). Factor inhibiting HIF (FIH-1) promotes renal cancer cell survival by protecting cells from HIF-1 α -mediated apoptosis. *Br J Cancer*, **104**, 1151-9.
- Kim TH, Hur EG, Kang SJ, et al (2011). NRF2 blockade suppresses colon tumor angiogenesis by inhibiting hypoxia-induced activation of HIF-1 α . *Cancer Res*, **71**, 2260-75.
- Kimura S, Kitadai Y, Tanaka S, et al (2004). Expression of hypoxia-inducible factor (HIF)-1 α is associated with vascular endothelial growth factor expression and tumour angiogenesis in human oesophageal squamous cell carcinoma. *Eur J Cancer*, **40**, 1904-12.
- Koukourakis MI, Giatromanolaki A, Skarlatos J, et al (2001). Hypoxia inducible factor (HIF-1 α and HIF-2 α) expression in early esophageal cancer and response to photodynamic therapy and radiotherapy. *Cancer Res*, **61**, 1830-2.
- Kurokawa T, Miyamoto M, Kato K, et al (2003). Overexpression of hypoxia-inducible-factor 1 α (HIF-1 α) in oesophageal squamous cell carcinoma correlates with lymph node metastasis and pathologic stage. *Br J Cancer*, **89**, 1042-7.
- Matsuyama T, Nakanishi K, Hayashi T, et al (2005). Expression of hypoxia-inducible factor-1 α in esophageal squamous cell carcinoma. *Cancer Sci*, **96**, 176-82.
- Rapisarda A, Shoemaker RH, Melillo G (2004). Targeting topoisomerase I to inhibit hypoxia inducible factor 1. *Cell Cycle*, **3**, 172-5.
- Seeber LM, Zweemer RP, Verheijen RH, et al (2010). Hypoxia-inducible factor-1 as a therapeutic target in endometrial cancer management. *Obstet Gynecol Int*, **2010**, 580971.
- Sithanandam G, Fornwald LW, Fields J, et al (2005). Inactivation of ErbB3 by siRNA promotes apoptosis and attenuates growth and invasiveness of human lung adenocarcinoma cell line A549. *Oncogene*, **24**, 1847-59.
- Takashima N, Ishiguro H, Kuwabara Y, et al (2008). Gene expression profiling of the response of esophageal carcinoma cells to cisplatin. *Dis Esophagus*, **21**, 230-5.
- Tzao C, Lee SC, Tung HJ, et al (2008). Expression of hypoxia-inducible factor (HIF)-1 α and vascular endothelial growth factor (VEGF)-D as outcome predictors in resected esophageal squamous cell carcinoma. *Dis Markers*, **25**, 141-8.
- Wang G, Ye Y, Yang X, et al (2011). Expression-based in silico screening of candidate therapeutic compounds for lung adenocarcinoma. *PLoS One*, **6**, e14573.
- Welsh S, Williams R, Kirkpatrick L, et al (2004). Antitumor activity and pharmacodynamic properties of PX-478, an inhibitor of hypoxia-inducible factor-1 α . *Mol Cancer Ther*, **3**, 233-44.
- Wu XA, Sun Y, Fan QX, et al (2007). Impact of RNA interference targeting hypoxia-inducible factor-1 α on chemosensitivity in esophageal squamous cell carcinoma cells under hypoxia. *Zhonghua Yi Xue Za Zhi*, **87**, 2640-4.
- Yonesaka K, Tamura K, Kurata T, et al (2006). Small interfering RNA targeting survivin sensitizes lung cancer cell with mutant p53 to adriamycin. *Int J Cancer*, **118**, 812-20.