

Alteration of Striatal Tetrahydrobiopterin in Iron-Induced Unilateral Model of Parkinson's Disease

Bijay Aryal¹, Jin-Koo Lee^{1,2}, Hak Rim Kim^{1,2}, and Hyung-Gun Kim^{1,2}

¹Department of Pharmacology, College of Medicine, ²Translational Research Center, Institute of Bio-Science Technology, Dankook University, Cheonan 330-714, Korea

It has been suggested that transition metal ions such as iron can produce oxidative injuries to nigrostriatal dopaminergic neurons, like Parkinson's disease (PD) and subsequent compensative increase of tetrahydrobiopterin (BH₄) during the disease progression induces the aggravation of dopaminergic neurodegeneration in striatum. It had been established that the direct administration of BH₄ into neuron would induce the neuronal toxicity *in vitro*. To elucidate a role of BH₄ in pathogenesis in the PD *in vivo*, we assessed the changes of dopamine (DA) and BH₄ at striatum in unilateral intranigral iron infused PD rat model. The ipsistriatal DA and BH₄ levels were significantly increased at 0.5 to 1 d and were continually depleting during 2 to 7 d after intranigral iron infusion. The turnover rate of BH₄ was higher than that of DA in early phase. However, the expression level of *GTP-cyclohydrolase I* mRNA in striatum was steadily increased after iron administration. These results suggest that the accumulation of intranigral iron leads to generation of oxidative stress which damage to dopaminergic neurons and causes increased release of BH₄ in the dopaminergic neuron. The degenerating dopaminergic neurons decrease the synthesis and release of both BH₄ and DA *in vivo* that are relevance to the progression of PD. Based on these data, we propose that the increase of BH₄ can deteriorate the disease progression in early phase of PD, and the inhibition of BH₄ increase could be a strategy for PD treatment.

Key Words: Dopamine, Ferrous citrate, GTP-cyclohydrolase I, Parkinson's disease, Tetrahydrobiopterin

INTRODUCTION

Parkinson's disease (PD) is a neurodegenerative disorder affecting 1% of population above the age of 65 [1] and is characterized by selective loss of dopaminergic neurons in the substantia nigra pars compacta (SNc). Degradation of the dopaminergic neurons located in the SNc and a subsequent loss of dopaminergic nerve terminals in the striatum are responsible for most of the movement disorders [2]. Although the underlying cause of dopaminergic cell death or process by which these cells degenerate in PD is not clearly understood but oxidative stress [3], mitochondrial dysfunction [4], apoptosis [5-7] and protein misfolding [8] are thought to play an important role in causing dopaminergic cell death and cellular degeneration.

Within the central nervous system, tetrahydrobiopterin

(BH₄) is an essential cofactor for dopamine (DA) and serotonin synthesis. The BH₄ is an obligatory cofactor for tyrosine hydroxylase (TH), the first and rate limiting enzyme in the dopamine synthesis and is produced selectively in monoaminergic neurons in the brain including the nigral dopaminergic neurons [9-11]. It has been reported that excess DA is potential source of oxidative stress [12] and is major contributing factor for nigrostriatal degeneration [13-15]. It has been demonstrated that BH₄ and DA together lead to generation of oxidative stress and render DA cells particularly vulnerable [16] and furthermore administration of BH₄ to animals leads to as selective degeneration of the DA system [17]. BH₄ itself generates oxidative stress by undergoing auto-oxidation [18-21] and elicits the formation of reactive quinonoid derivatives of DA and protein as well as lipid peroxidation in DA cells [22]. The amount of BH₄ synthesizing enzyme, GTP cyclohydrolase I (*GTP-CH I*), is lowest, suggesting that the level of BH₄ in this area might be normally subtoxic [23,24]. However, it is possible that conditions that lead to aberrant increase in BH₄ and DA might elicit oxidative damage at SNc.

It has been reported that transition metal ions like, iron

Received November 20, 2013, Revised January 19, 2014,
Accepted February 5, 2014

Corresponding to: Hyung-Gun Kim, Department of Pharmacology, College of Medicine, Dankook University, 119 Dandaero, Dongnam-gu, Cheonan 330-714, Korea. (Tel) 82-41-550-3867, (Fax) 82-41-551-3866, (E-mail) hgkimm@dankook.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABBREVIATIONS: PD, Parkinson's disease; SNc, substantia nigra pars compacta; BH₄, tetrahydrobiopterin; DA, dopamine; TH, tyrosine hydroxylase; *GTP-CH I*, GTP cyclohydrolase I; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; LC-MS/MS, liquid chromatography with triple quadrupole tandem mass spectrometer.

has dopaminergic neurotoxin properties similar to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) because of its capacity to inducing brain injury through the generation of reactive oxygen species and the acceleration of DA auto-oxidation [25]. The iron related striatal DA depletion or turnover via DA auto-oxidation or oxidative stress on PD animal models have been already reported [26] however the iron related striatal BH₄ depletion or turnover via oxidative stress has not been studied yet. Hence, the present study aims to investigate the effect of intranigral infusion of ferrous citrate on DA and BH₄ turnover in rat nigrostriatal system in the progress of PD. We also analyzed *GTP-CH I* mRNA expression in rat nigrostriatal system in parallel.

METHODS

Animal handling and care

All the experimental protocols involving animals were reviewed and approved by the institutional animal care and use committee (IACUC) of Dankook University. Male Sprague Dawley rats (250~300 g) were purchased from Daehan Biolink, Korea and upon the arrival, the animals were housed three per cage for two weeks at room temperature of 22±2°C and relative humidity 60~70% on a 12-h light-dark cycle with adequate supply of standard rodent food and water *ad libitum*.

Unilateral destruction of nigrostriatal dopaminergic neuron

Ferrous citrate was diluted in sterile 0.9% NaCl solution to produce the concentration of 2 mM. The solution was filtered through 0.2 μm Minisart® filter unit and was used for intranigral administration. On the day of surgery, rats were anaesthetized with intraperitoneally (i.p.) injection of tiletamine hydrochloride (125 mg/kg) and zolazepam hydrochloride (125 mg/kg) and positioned on stereotaxic instrument (Kopf Instruments, Tujunga, CA, USA). The 4.0 μL of ferrous citrate solution (2 mM) was infused through microdialysis probe with glued outlet and tear off membrane (CMA12®, CMA/Microdialysis AB, Stockholm, Sweden) at the flow rate of 0.8 μL/min to the right SNC using stereotaxic coordinates; AP: 3.2 mm, L: 2.1 mm and DV: 2.0 mm relative to bregma by the mouse brain atlas [27]. The injections were delivered over the period of 5 min and probes were kept *in situ* for an additional 5 min after each injection to avoid the reflux along the injection tract. The sham controls were injected with 4.0 μL of 0.9% NaCl. After surgery, animals were decapitated at different time intervals 0 h, 12 h, 24 h, 48 h, 96 h and 1 wk (n=6) using guillotine cutter and brain was quickly removed. The brain was dissected on an ice plate to separate right (ipsilateral) and left (contralateral) striatum. Both striatum were safely stored at -70°C for biochemical analysis. Striatum was weighted and homogenized in acetonitrile (1 : 10, w/w) with 5 μL of IS (250 ng/mL). After homogenizing thoroughly, the DA and BH₄ from striatum were further extracted by sonication for 60 sec. The homogenates of brain tissues were centrifuged by 12,000 rpm for 10 min at 4°C. Supernatants were transferred to 96-well plate and then injected onto the LC-MS/MS system by auto-sampler for subsequent analysis.

LC-MS/MS determination of DA and BH₄

The modified method was used for simultaneous determination of DA and BH₄ in brain sample by using liquid chromatography with triple quadrupole tandem mass spectrometer (LC-MS/MS) [28]. Briefly striatum was accurately weighed and mixed with acetonitrile containing 200 μg/ml dithiothreitol and 2 μg/ml cimetidine (internal standard). The homogenized mixture was vortexed for 30 sec and centrifuged (8,000 rpm, 10 min) for extraction of DA and BH₄ from striatum. The clear supernatant was collected in 100 μL of microtube and was dried under gentle stream of nitrogen at 45°C. The dried extract was reconstituted with 100 μL of 50% methanol containing 0.1% formic acid and 10 μL was injected into LC-MS/MS system. The sample was eluted with mobile phase and was analyzed under the multiple reaction monitoring (MRM) mode: DA (m/z 154 → 177, CE=10 eV), BH₄ (m/z 242.2 → 166.0, CE=17 eV), cimetidine (m/z 253.1 → 159.0, CE=14 eV). The liquid chromatographic system was a Accela (Thermo Fisher Scientific Inc., Waltham, MA, USA) system equipped with a solvent delivery module, Nanospace SI-2 3133 (Shiseido Inc., Japan) with an autosampler, and connected to a Discovery Max (Thermo Fisher Scientific Inc.) quadrupole tandem mass spectrometer equipped with electrospray ionization. System control and data analyses were carried out with Xcalibur (Thermo Fisher Scientific Inc.). Chromatographic separation was achieved using a hydrophilic interaction chromatography HPLC column (Luna 3u HILIC 200A, Phenomenex, Torrance, CA, USA) protected by a guard column (Phenomenex C₁₈, 4×2 mm, Phenomenex). A column oven (Nanospace SI-2 3004, Shiseido, Japan) was used online. The mobile phase of 0.1% formic acid-acetonitrile/water (75 : 25, v/v) was run at a flow rate of 300 μL/min. The electrospray ionization (ESI) mass spectrometer was operated in the positive ion mode.

Total RNA isolation and reverse transcription

The brain tissues were homogenized in TRIzol® reagent (Invitrogen). Total RNA was extracted from the cells according to the manufacturer's suggested protocol. Total RNA (2 μg) was treated with 1 U DNase I (Promega) for 15 min at RT in 18 μL of volume containing 1X PCR buffer and MgCl₂ (2 mM). Then it was inactivated by incubation with 2 μL of EDTA (25 mM) at 65°C for 15 min. Reverse transcriptase reactions were carried out using MuLV Reverse Transcriptase (Promega) according to the manufacturer's protocol. Each reaction tube contained 2 μg of total in a volume of 25 μL containing 5 μL of MuLV 5X RT buffer, 1 μg of oligo-d(T) (promega), 2.5 μL of dNTP Mixture (Promega), 40 U of RNasin (Promega), 20 U of MuLV Reverse Transcriptase and nuclease-free water to volume. Reverse transcriptase reactions were carried out in a DNA Thermal Cycler 480 (Perkin Elmer, Branchburg, NJ, U.S.A.) at 25°C for 20 min, 42°C for 60 min and 95°C for 10 min. The cDNA was then stored at -20°C until use.

Quantitative real time PCR

Real-time PCR for the analysis of *GTP-CH I* mRNA levels were performed in a Rotor-Gene 3000 (Corbett life sciences, RG-3000, USA). QuantiTect SYBR Green PCR kit and primers (rat *GTP-CH I* and *GAPDH* genes) were purchased

from Qiagen. The reaction mixture consisted of 2 μ L of cDNA template, 10 μ L of SYBR Green PCR master mix and 10 pmol of primers in total volume of 20 μ L. The cDNA was denatured at 95°C for 5 min followed by 40 cycles of PCR at 95°C for 5 sec, 55°C for 10 sec and 72°C for 15 sec. Data acquisition and analysis of real-time PCR were performed using the Rotor-Gene 3000 series software. Delta-delta Ct method was used to calculate the relative quantita-

tion of *GTP-CH 1* normalized with *GAPDH* level in each individual sample.

Statistical Analysis

All values shown in the figures are expressed as the mean \pm SEM obtained from at least three independent experiments. Statistical analysis was carried out one-way analysis of variance (ANOVA) with Tukey's post-hoc test (Fig. 2~4) using GraphPad Prism (GraphPad Software, San Diego, CA, USA). Statistically significant differences between groups were assumed when $p < 0.05$.

RESULTS

Histological verification of unilateral intranigral iron infusion

As the preliminary experiment, upon completion of the microinfusion of ferrous citrate into SN, the animals were anaesthetized with i.p. injection of tiletamine hydrochloride (125 mg/kg) and zolazepam hydrochloride (125 mg/kg) and transcardially perfused with 10% formalin (neutral buffered, pH 7.4). The brains were separated and immersed in the fixative for overnight and then brain tissues were cryoprotected after soaking in a series of sucrose solution (10%, 20%, and 30%) at 4°C until they sank in liquid nitrogen. Serial coronal section, 40 μ m were cut with freezing, sliding microtome and stained with cresyl violet. The image shows that injected microdialysis probe tips were located in the SN which was stained with iron (Fig. 1).

Observation of behavioral response after unilateral intranigral iron infusion

After unilateral intranigral infusion of ferrous citrate, animals showed spontaneous contralateral circling behavior. Although spontaneous circling behavior gradually reduced after lesion, it was exaggerated by the application




Fig. 1. Histological verification of unilateral intranigral iron infusion. Coronal section image after unilateral intranigral iron infusion (A). Arrow indicate injected site of microdialysis probe tip. Crystal violet staining in ferrous citrate infused SN (ipsilateral SN) (B). SNC, substantia nigra pars compacta; SNR, substantia nigra pars reticularis.




Fig. 2. Alterations of ipsilateral DA levels in intranigral iron infused rats. Wet tissue was isolated from both ipsilateral and contralateral of intranigral unilateral saline (4.0 μ L) or ferrous citrate (4.0 μ L, 2 mM) infused rats at 0, 0.5, 1, 2, 4, and 7 days after challenge. DA level was measured by LC-MS/MS. The levels of DA (ng/wet tissue weight g) convert to % of ipsilateral / contralateral level. Values are mean \pm SEM (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; compared to control group, $n=6$).




Fig. 3. Alterations of ipsilateral BH₄ levels in intranigral iron infused rats. Wet tissue was isolated from both ipsilateral and contralateral of intranigral unilateral saline (4.0 μ L) or ferrous citrate (4.0 μ L, 2 mM) infused rats at 0, 0.5, 1, 2, 4, and 7 days after challenge. BH₄ level was measured by LC-MS/MS. The levels of BH₄ (ng/wet tissue weight g) convert to % of ipsilateral / contralateral level. Values are mean \pm SEM (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; compared to control group, $n=6$).




Fig. 4. Changes of ipsilateral *GTP-CH I* mRNA levels in intranigral iron infused rats. Total RNA was isolated from both ipsilateral and contralateral striatum of intranigral unilateral saline (4.0 μ L) or ferrous citrate (4.0 μ L, 2 mM) infused rats at 0, 0.5, 1, 2, 4, and 7 days after challenge. Gene expression of *GTP-CH I* was measured by quantitative real-time PCR using Rotor-Gene 3000. The mRNA level of *GTP-CH I* (ipsilateral / contralateral) was normalized with GAPDH gene expression. Data were obtained from triplicated PCR reactions, and values are mean \pm SEM (* p < 0.05, ** p < 0.01, *** p < 0.001; compared to control group, n =6).

of painful stimulus. The contralateral circling movement was increased after pinch application by pincette on the left hindlimb or drop of hot water (70°C) by Pasteur pipette on left ear (data not shown). It was suggested that the circling movement after surgery was due to imbalance of nigrostriatal dopamine in lesioned and intact hemispheres, and contralateral circling movement is the key behavioral phenotype of the unilateral lesioned PD rodent model [29]. There were no significance between contralateral circling behavior and DA level (data not shown).

Alteration of striatal DA level

Intranigral infusion of ferrous citrate caused changing DA releases in the striatum via nigrostriatal dopaminergic system. The average concentration of DA in the contralateral striatum (n =12) was 4328.5 \pm 26.4 (ng/wet tissue weight g). The percentage ratio of DA concentration in the ipsilateral striatum (n =6) after stereotaxic iron infusion at different time intervals (0, 0.5, 1, 2, 4, and 7 days) were measured. The maximum ipsilateral DA level was 200.9 \pm 10.7% at 1 d and ipsilateral DA was severely depleted to 2.7 \pm 1.1% at 4 d after intranigral infusion of 4 μ L of 2 mM ferrous citrate (Fig. 2) The DA levels in ipsilateral striatum by the intranigral infusion of 4 μ L of 0.9% NaCl (vehicle, n =4) with respect to percentage of contralateral striatum at 0 and 7 d were 97.0 \pm 3.7% and 98.9 \pm 6.7%, respectively. There were significant differences of ipsilateral DA concentration between 0 d (before injection) and 0.5, 1, 2, 4, and 7 days after intranigral ferrous citrate administration (Fig. 2). These data showed that intranigral iron accumulation caused initial increase in DA levels in nigrostriatal system which induced oxidative stress, preceded chronic decrease in the striatal DA levels such as the progress of PD.

Alteration of striatal BH₄ level

Intranigral infusion of ferrous citrate caused changing

BH₄ release in the striatum by nigrostriatal dopaminergic system in a similar pattern of DA release. The average concentration of BH₄ in the contralateral striatum (n =12) was 192.1 \pm 19.0 (ng/wet tissue weight g). The percentage ratio of BH₄ concentration in the ipsilateral striatum (n =6) after stereotaxic iron infusion at different time intervals from 0 to 7 d were measured. The maximum ipsilateral BH₄ level was 244.7 \pm 2.0% at 1 d and ipsilateral BH₄ was severely depleted to 4.5 \pm 3.5% at 4 d after intranigral iron infusion (Fig. 3) The BH₄ levels in ipsilateral striatum by the intranigral infusion of vehicle (n =4) with respect to percentage of contralateral striatum at 0 and 7 d were 94.9 \pm 12.3% and 96.3 \pm 11.7%, respectively. There were significant differences of ipsilateral BH₄ level between 0 d (before injection) and 0.5, 1, 2, 4, and 7 days after intranigral ferrous citrate administration (Fig. 3). These data showed that intranigral iron accumulation caused initial increase in BH₄ levels in nigrostriatal dopaminergic neurons, which preceded chronic decrease in the striatal BH₄ levels. The result showed that the turnover of BH₄ and DA were significantly correlated (r =0.968, p <0.05) after iron infusion. However, striatal BH₄ concentration increased further than DA concentration until 1 d but interestingly, decreased quickly after 1 d compared with striatal DA turnover in rat nigrostriatal system. It is suggested that the BH₄ concentration changes more dynamically than the DA concentration in striatal system (Fig. 2 and 3).

Changes of GTP-CH I mRNA expression

Intranigral infusion of ferrous citrate induced persistent increase of *GTP-CH I* mRNA expression in the striatum. The percentage ratio of quantitative real time PCR values for *GTP-CH I* mRNA expression on the ipsilateral striatum (injected side, n =6) to the contralateral side after stereotaxic iron infusion at different time intervals from 0 to 7 d were increased gradually up to 277.4 \pm 4.4% at 7 d after ferrous citrate administration (Fig. 4). The *GTP-CH I* mRNA expression by the intranigral infusion of vehicle with respect to percentage of ipsi-/contra-lateral striatum at 0 and 7 d were 101.2 \pm 3.2% and 106.2 \pm 9.3%, respectively. There were significant differences of *GTP-CH I* mRNA expression between 0 d and 0.5, 1, 2, 4, and 7 days after intranigral ferrous citrate infusion (Fig. 4). These data showed that intranigral iron accumulation induced *GTP-CH I* mRNA expression immediately and the induction of gene expression was sustained at least 7 d in striatum, suggesting increased *GTP-CH I* mRNA expression might be a compensatory process of BH₄ which depleted by the degenerative progress of nigrostriatal dopaminergic neurons.

DISCUSSION

It has been known that trace metal, iron, was accumulated in brain with aging and induced oxidative stress in the basal ganglia, especially in the A9 dopaminergic neurons of SNC area [29]. Interestingly, non-heme iron was increased further in the SN of patients with PD [30-32]. In the present study, we had microinfused ferrous citrate, a small molecular weight iron complex, into the SNC area unilaterally as the pathophysiological animal model mimicking PD to study the pathogenic path of the PD. We found that iron caused a chronic depletion of striatal DA and BH₄ levels, which was preceded by initial increase in striatal

DA and BH₄ concentration. In line with our findings, BH₄ levels are decreased at 24 h in MPP⁺ (1-methyl-4-phenylpyridinium)-induced human dopaminergic model of PD [33]. Previous our findings show that MPTP induce the reduction of DA but not BH₄ level in striatum and SN [34]. Intranigral iron infusion induced nigral DA and BH₄ release into striatum peaked during 0.5 and 1 d after the stereotaxic procedure which suggested that initial increased synthesis and/or release of nigral DA and BH₄ induced by iron accumulation augmented the oxidative damage to nigrostriatal dopaminergic neurons, and damaged nigrostriatal dopaminergic neurons represented reduced synthesizing capacity of DA and BH₄ as the biochemical phenotype in PD. Therefore, this intranigral iron infusion is a quite reliable model for PD because it showed both biochemical and behavioral phenotypes of PD with slow progress even similar neurochemical and behavioral changes have been documented following infusion of dopaminergic neurotoxin MPTP and 6-hydroxydopamine in the nigrostriatal dopaminergic system [34-36].

It has been previously proposed that oxidative injury in PD may be caused by oxidants generated during enhanced DA turnover or iron-catalyzed DA auto-oxidation [37,38]. Iron induced hydroxyl radical generation and lipid peroxidation may be caused, at least in part, by sustained DA turnover which was observed during first two days after iron infusion. Earlier reports indicated that iron induced striatal DA depletion is linked to the loss of tyrosine hydroxylase-positive neurons in the SN. The present study demonstrates that the degree of the nigrostriatal degeneration is correlated with, not only DA turnover, but also chronic depletion of striatal BH₄. Several studies showed that both hydroxyl radicals and iron have been implicated in the initiation and propagation of chain reaction of lipid peroxidation [30-32]. It has been well documented that iron caused sustained lipid peroxidation which induced cell membrane damage, calcium overload and eventually lead to cell death when the oxidative stress overwhelms the cellular integrity [31]. Elevated non-heme iron is routinely found in the SNC in PD patients [32]. As suggested by Kim et al. [39], it is possible that an aberrant increase in BH₄ and DA might elicit oxidative damage in the SNC, and resulting structural damage to nigrostriatal pathway.

BH₄ auto-oxidation could produce hydrogen peroxide and superoxide radical [18-21]. In addition, DA, with its synthesis facilitated by BH₄, in turn undergoes auto-oxidation to form a reactive quinone species [18,22,24]. Indeed, DA derived quinone production is elevated in which BH₄ provides its auto-oxidation products [18]. As both BH₄ and DA are inevitably produced within the same cells for DA neurotransmission, the DA cells would tend to be more vulnerable especially under intranigral iron infusion conditions that result in overproduction of BH₄ and DA.

All the monoaminergic neurons and terminals in the CNS produce BH₄ and DA cells are vulnerable to high level of BH₄, most probably due to the higher rate of oxidation of DA compared with other catecholamine or serotonin [18,19]. Under, the normal conditions, the level of *GTP-CH I* expression in the nigrostriatal system is lower than in the other monoaminergic cells [23,24]. In addition, DA is normally sequestered in the storage vesicles by the action of vesicular monoamine transporter (VMAT) after synthesis in the cytosol, limiting its encounter with BH₄ [23]. Therefore, an aberrant condition such as intranigral iron infusion, would lead to initial overproduction of BH₄ and DA

as the consequence of oxidative damage and then gradually depress the production of BH₄ and DA, in which *GTP-CH I* levels were increased after 24 h with compared to sham control (vehicle infused group).

This is the first study for demonstrating an overproduction of BH₄ and DA actually occur *in vivo* in PD animal model. Iron accumulation in SN elevated BH₄ and its synthesizing enzymes, *GTP-CH I* and DA. This result provides a strong support for current view in progression of PD that excessive DA is a contributing factor for nigrostriatal degeneration. In addition, we demonstrate that intranigral infusion of iron could generate a PD animal model for studying pathophysiology and for developing preventive method of PD progress *in vivo*. In the present study we showed iron accumulation in SN causes increases in production of both BH₄ and DA and this concomitant elevation induces degeneration of the nigrostriatal pathway. As BH₄ can increase oxidative damage by auto-oxidation as well as by enhancing synthesis of DA, whose oxidation is in turn facilitated in the presence of BH₄, it is possible that the iron accumulation in the dopaminergic neuron can increase BH₄ levels, which might be a primary etiologic factor of PD in aged brain.

ACKNOWLEDGEMENTS

The present research was conducted by the research fund of Dankook university in 2012.

REFERENCES

1. Zhang Y, Dawson VL, Dawson TM. Oxidative stress and genetics in the pathogenesis of Parkinson's disease. *Neurobiol Dis.* 2000;7:240-250.
2. Hwang O, Choi HJ, Park SY. Up-regulation of GTP cyclohydrolase I and tetrahydrobiopterin by calcium influx. *Neuroreport.* 1999;10:3611-3614.
3. Beal MF. Mitochondria, oxidative damage, and inflammation in Parkinson's disease. *Ann N Y Acad Sci.* 2003;991:120-131.
4. Greenamyre JT, Sherer TB, Betarbet R, Panov AV. Complex I and Parkinson's disease. *IUBMB Life.* 2001;52:135-141.
5. Anglade P, Vyas S, Javoy-Agid F, Herrero MT, Michel PP, Marquez J, Mouatt-Prigent A, Ruberg M, Hirsch EC, Agid Y. Apoptosis and autophagy in nigral neurons of patients with Parkinson's disease. *Histol Histopathol.* 1997;12:25-31.
6. Kingsbury AE, Mardsen CD, Foster OJ. DNA fragmentation in human substantia nigra: apoptosis or perimortem effect? *Mov Disord.* 1998;13:877-884.
7. Tatton WG, Chalmers-Redman R, Brown D, Tatton N. Apoptosis in Parkinson's disease: signals for neuronal degradation. *Ann Neurol.* 2003;53 Suppl 3:S61-70; discussion S70-72.
8. McNaught KS, Olanow CW. Proteolytic stress: a unifying concept for the etiopathogenesis of Parkinson's disease. *Ann Neurol.* 2003;53 Suppl 3:S73-84; discussion S84-86.
9. Hwang O, Baker H, Gross S, Joh TH. Localization of GTP cyclohydrolase in monoaminergic but not nitric oxide-producing cells. *Synapse.* 1998;28:140-153.
10. Kaufman S. New tetrahydrobiopterin-dependent systems. *Annu Rev Nutr.* 1993;13:261-286.
11. Nagatsu I, Ichinose H, Sakai M, Titani K, Suzuki M, Nagatsu T. Immunocytochemical localization of GTP cyclohydrolase I in the brain, adrenal gland, and liver of mice. *J Neural Transm Gen Sect.* 1995;102:175-188.
12. Offen D, Ziv I, Sternin H, Melamed E, Hochman A. Prevention of dopamine-induced cell death by thiol antioxidants: possible implications for treatment of Parkinson's disease. *Exp Neurol.*

- 1996;141:32-39.
13. **Asanuma M, Miyazaki I, Ogawa N.** Dopamine- or L-DOPA-induced neurotoxicity: the role of dopamine quinone formation and tyrosinase in a model of Parkinson's disease. *Neurotox Res.* 2003;5:165-176.
 14. **Hastings TG, Zigmond MJ.** Loss of dopaminergic neurons in parkinsonism: possible role of reactive dopamine metabolites. *J Neural Transm Suppl.* 1997;49:103-110.
 15. **Weingarten P, Zhou QY.** Protection of intracellular dopamine cytotoxicity by dopamine disposition and metabolism factors. *J Neurochem.* 2001;77:776-785.
 16. **Choi HJ, Jang YJ, Kim HJ, Hwang O.** Tetrahydrobiopterin is released from and causes preferential death of catecholaminergic cells by oxidative stress. *Mol Pharmacol.* 2000;58:633-640.
 17. **Choi HJ, Kim SW, Lee SY, Hwang O.** Dopamine-dependent cytotoxicity of tetrahydrobiopterin: a possible mechanism for selective neurodegeneration in Parkinson's disease. *J Neurochem.* 2003;86:143-152.
 18. **Davis MD, Kaufman S.** Products of the tyrosine-dependent oxidation of tetrahydrobiopterin by rat liver phenylalanine hydroxylase. *Arch Biochem Biophys.* 1993;304:9-16.
 19. **Davis MD, Kaufman S, Milstien S.** The auto-oxidation of tetrahydrobiopterin. *Eur J Biochem.* 1988;173:345-351.
 20. **Kirsch M, Korth HG, Stenert V, Sustmann R, de Groot H.** The autoxidation of tetrahydrobiopterin revisited. Proof of superoxide formation from reaction of tetrahydrobiopterin with molecular oxygen. *J Biol Chem.* 2003;278:24481-24490.
 21. **Thoeni G, Werner ER, Werner-Felmayer G.** Tetrahydropteridines suppress gene expression and induce apoptosis of activated RAW264.7 cells via formation of hydrogen peroxide. *Free Radic Biol Med.* 2004;37:375-385.
 22. **Choi HJ, Lee SY, Cho Y, Hwang O.** Inhibition of vesicular monoamine transporter enhances vulnerability of dopaminergic cells: relevance to Parkinson's disease. *Neurochem Int.* 2005;46:329-335.
 23. **Double KL, Ben-Shachar D, Youdim MB, Zecca L, Riederer P, Gerlach M.** Influence of neuromelanin on oxidative pathways within the human substantia nigra. *Neurotoxicol Teratol.* 2002;24:621-628.
 24. **Lentz SI, Kapatos G.** Tetrahydrobiopterin biosynthesis in the rat brain: heterogeneity of GTP cyclohydrolase I mRNA expression in monoamine-containing neurons. *Neurochem Int.* 1996;28:569-582.
 25. **Keane PC, Kurzawa M, Blain PG, Morris CM.** Mitochondrial dysfunction in Parkinson's disease. *Parkinsons Dis.* 2011;2011:716871.
 26. **Eşrefoğlu M.** Cell injury and death: Oxidative stress and antioxidant defense system: Review. *Turkiye Klinikleri J Med Sci.* 2009;29:1660-1676.
 27. **Paxinos G, Franklin K.** The mouse brain in stereotaxic coordinates. Compact: The Coronal Plates and Diagrams. Allen Institute: Academic Press; 2008.
 28. **Zhao Y, Cao J, Chen YS, Zhu Y, Patrick C, Chien B, Cheng A, Foehr ED.** Detection of tetrahydrobiopterin by LC-MS/MS in plasma from multiple species. *Bioanalysis.* 2009;1:895-903.
 29. **Mokřý J.** Experimental models and behavioural tests used in the study of Parkinson's disease. *Physiol Res.* 1995;44:143-150.
 30. **Dexter DT, Carayon A, Javoy-Agid F, Agid Y, Wells FR, Daniel SE, Lees AJ, Jenner P, Marsden CD.** Alterations in the levels of iron, ferritin and other trace metals in Parkinson's disease and other neurodegenerative diseases affecting the basal ganglia. *Brain.* 1991;114:1953-1975.
 31. **Gerlach M, Ben-Shachar D, Riederer P, Youdim MB.** Altered brain metabolism of iron as a cause of neurodegenerative diseases? *J Neurochem.* 1994;63:793-807.
 32. **Morris CM, Keith AB, Edwardson JA, Pullen RG.** Uptake and distribution of iron and transferrin in the adult rat brain. *J Neurochem.* 1992;59:300-306.
 33. **Ryan BJ, Lourenço-Venda LL, Crabtree MJ, Hale AB, Channon KM, Wade-Martins R.** α -Synuclein and mitochondrial bioenergetics regulate tetrahydrobiopterin levels in a human dopaminergic model of Parkinson disease. *Free Radic Biol Med.* 2013;67C:58-68.
 34. **Lee KS, Lee JK, Kim HG, Kim HR.** Differential Effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine on Motor Behavior and Dopamine Levels at Brain Regions in Three Different Mouse Strains. *Korean J Physiol Pharmacol.* 2013;17:89-97.
 35. **Lotharius J, O'Malley KL.** The parkinsonism-inducing drug 1-methyl-4-phenylpyridinium triggers intracellular dopamine oxidation. A novel mechanism of toxicity. *J Biol Chem.* 2000;275:38581-38588.
 36. **Choe MA, Koo BS, An GJ, Jeon S.** Effects of Treadmill Exercise on the Recovery of Dopaminergic Neuron Loss and Muscle Atrophy in the 6-OHDA Lesioned Parkinson's Disease Rat Model. *Korean J Physiol Pharmacol.* 2012;16:305-312.
 37. **Fahn S, Cohen G.** The oxidant stress hypothesis in Parkinson's disease: evidence supporting it. *Ann Neurol.* 1992;32:804-812.
 38. **Jenner P, Olanow CW.** Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology.* 1996;47(6 Suppl 3):S161-170.
 39. **Kim ST, Choi JH, Chang JW, Kim SW, Hwang O.** Immobilization stress causes increases in tetrahydrobiopterin, dopamine, and neuromelanin and oxidative damage in the nigrostriatal system. *J Neurochem.* 2005;95:89-98.