

Ursolic Acid-Induced Elevation of Serum Irisin Augments Muscle Strength During Resistance Training in Men

Hyun Seok Bang^{1*}, Dae Yun Seo^{2*}, Yong Min Chung³, Kyoung-Mo Oh⁴, Jung Jun Park⁵, Figueroa Arturo⁶, Seung-Hun Jeong², Nari Kim², and Jin Han²

¹Division of Humanities and Social Science, POSTECH, Pohang 790-784, ²Department of Physiology, College of Medicine, Cardiovascular and Metabolic Disease Center, Inje University, Busan 614-735, ³Department of Physical Education, Tongmyong University, Busan 608-711, ⁴Department of Physical Education, Pukyong University, Busan 608-737, ⁵Division of Sport Science, Pusan National University, Busan 609-735, Korea, ⁶Department of Nutrition, Food and Exercise Sciences, Florida State University, Tallahassee FL 32306, USA

Ursolic acid (UA), a type of pentacyclic triterpenoid carboxylic acid purified from natural plants, can promote skeletal muscle development. We measured the effect of resistance training (RT) with/without UA on skeletal muscle development and related factors in men. Sixteen healthy male participants (age, 29.37±5.14 years; body mass index=27.13±2.16 kg/m²) were randomly assigned to RT (n=7) or RT with UA (RT+UA, n=9) groups. Both groups completed 8 weeks of intervention consisting of 5 sets of 26 exercises, with 10~15 repetitions at 60~80% of 1 repetition maximum and a 60~90-s rest interval between sets, performed 6 times/week. UA or placebo was orally ingested as 1 capsule 3 times/day for 8 weeks. The following factors were measured pre-and post-intervention: body composition, insulin, insulin-like growth factor-1 (IGF-1), irisin, and skeletal muscle strength. Body fat percentage was significantly decreased (p<0.001) in the RT+UA group, despite body weight, body mass index, lean body mass, glucose, and insulin levels remaining unchanged. IGF-1 and irisin were significantly increased compared with baseline levels in the RT+UA group (p<0.05). Maximal right and left extension (p<0.01), right flexion (p<0.05), and left flexion (p<0.001) were significantly increased compared with baseline levels in the RT+UA group. These findings suggest that UA-induced elevation of serum irisin may be useful as an agent for the enhancement of skeletal muscle strength during RT.

Key Words: IGF-1, Irisin, Muscle strength, Resistance training, Ursolic acid

INTRODUCTION

Resistance training (RT) precipitates an anabolic environment by inducing both hormonal and molecular variations in skeletal muscle. RT contribute to enhancement of insulin, insulin-like growth factor-1 (IGF-1), and myokines, which are well known as anabolic hormones [1-3]. Specially, RT-induced elevation in IGF-1, which can regulate muscle hypertrophy and muscle strength, has been strongly associated with myokine levels [4]. Numerous studies indicate that the novel myokine irisin, which is a cleaved and se-

creted form of fibronectin III domain-containing 5 found in organs including brain, heart, adipose tissue, and skeletal muscle, provokes browning of white adipose tissue a process controlled by uncoupling protein 1 and peroxisome proliferator-activated receptor-gamma co-activator-1 alpha [5]. Some authors have suggested that exercise-mediated increase in circulating irisin was strongly related with improvement of body fat mass, insulin resistance, and IGF-1 [6-10]. Furthermore, a study by Bostrom et al. [11] suggested that the plasma level of circulating irisin increased after 10 weeks of endurance training. In contrast, Norheim et al. [12] and Moraes et al. [13] found a decrease in the plasma concentration of irisin after 12 weeks of training and 6 months of resistance exercise, respectively. Therefore, the precise role of irisin in exercise-related physiological change is currently controversial [14].

The benefits of nutritional strategies to increase muscle strength have been widely investigated for potential improvements to health outcomes in humans [15-19]. In this context, supplementation with phytochemical compounds has received significant attention for its potential applications to resistance training. Ursolic acid (UA) is an isomer

Received August 19, 2014, Revised September 19, 2014,
Accepted September 25, 2014

Corresponding to: Jin Han, Department of Physiology, College of Medicine, Cardiovascular and Metabolic Disease Center, Inje University, Bokgiro 75, Busangin-gu, Busan 614-735, Korea.
(Tel) 82-51-890-6727, (Fax) 82-51-891-8748, (Email) phyhanj@inje.ac.kr

*Both authors contributed equally to this work.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABBREVIATIONS: UA, ursolic acid; RT, resistance training; IGF-1, insulin-like growth factor-1; 1RM, 1 repetition maximum.

of oleanolic acid and a lipophilic triterpenoid compound primarily found in apples and many herbs [20]. Recently, UA has been demonstrated to have important biological roles including anti-fat accumulation [21], anti-insulin resistance via IGF-1 [6], anti-muscle atrophy [22], anti-cancer [23], anti-oxidation [24], and anti-inflammatory effects [25]. Above all, the underlying mechanism of UA is the enhancement or maintenance of skeletal muscle mass and function in vivo and vitro [22,25-27]. However, it is not clear whether increase in skeletal muscle mass is associated with signaling via the same IGF-1-irisin-pathway found to promote skeletal muscle strength in existing clinical studies. To date, several studies have assessed the impact of UA supplementation on muscle atrophy or obesity, but no reports have elucidated the biological effect of UA supplementation on human subjects involved in RT. For that purpose, we determined differences in body composition, fasting glucose, insulin, IGF-1, irisin levels, and muscle strength following RT with or without concomitant UA supplementation. We hypothesized that UA-induced elevation of serum irisin would promote body fat loss and improvements in skeletal muscle development during RT in men.

METHODS

Study design

The participants of this study performed an exercise program that included supervised RT with or without UA supplementation for 8 weeks. Body composition, blood parameters and muscular strength were determined at baseline and after this program. All participants signed written informed consent approved by the Pusan National University Institutional Research Board (PNU IRB/2014_42_HR).

Subjects

Twenty-four Korean men with over 3 years of RT experience were recruited from Dong Myoung University by newspapers and e-mail for this study. All respondents completed a family history and detailed health history questionnaires. Respondents were excluded from this study (n=8), if they met exclusion criteria including diagnosis of chronic disease including cardiovascular disease, hypertension, cancer, kidney dysfunction, or musculoskeletal dysfunction during the previous 3 months. Sixteen healthy participants (age: 29.37 ±5.14 years) were ultimately selected to participate in this study. All eligible participants were nonsmokers and had not taken medications or steroid therapy within 1 year prior to or during this study period. Participants were randomly divided into the following interventions: RT with placebo supplementation (RT; age: 28.71±5.76 years, n=7) and RT with UA supplementation (RT+UA; age: 29.88±4.91 years, n=9). Before the investigation, participants stopped consumption of other ergogenic aids immediately and were subjected to a 5-week washout period.

Resistance training

Participants performed the RT program for 8 weeks using free weights and machines under the supervision of a trained instructor, who directed stretching exercises as a warm up to prevent soft tissue injuries and ensure subject safety. The accuracy of a prediction equation to estimate

1 repetition maximum intensity (1RM) until executed to fatigue was evaluated at each exercise session before starting the program [27]. This RT program consisted of a total of 26 exercise types, with 13 upper-body (chin-up, lateral pull-down, bent over/up-right row, bench press, fly, cable cross-over, shoulder press, lateral raise, biceps curl, concentration curl, barbell elbow extension, and kick back) and 13 lower-body training exercises (squat, leg extension, leg press, sissy squat, leg curl, dead lift, stiff leg dead lift, lunges, back extension, seated/standing calf raise, sit-up, and abdominal crunch) to strengthen the extremities. Participants performed 5 sets of 10~15 repetitions at 60~80% of the 1RM with a 60~90-s inter-set rest, 6 times/week for 8 weeks.

Ursolic acid supplementation

The placebo group was provided with 3 capsules daily containing 450 mg of guar gum, and the UA group received 3 capsules daily containing 450 mg of UA from rosemary leaf extract (Labrada, Houston, Texas, USA). The capsules had the same color, shape, and size, and participants took 1 capsule (150 mg) after each of 3 meals. We monitored the participants by questionnaire via e-mail, cell phone, and visits to the laboratory to confirm compliance with supplementation throughout the study.

Anthropometric and biochemical parameters

Blood samples were collected before and after the 8-week intervention. Body weight and height were measured with the participants wearing light clothes and no shoes. Body composition including body weight, body fat percentage and lean body mass was measured before and after the 8-week intervention using a multi-frequency electrical impedance analyzer (X-scan Plus II, Jawon Medical, Seoul, Korea). Body mass index was calculated as body weight (kg) divided by the square of height (m). Blood samples were collected from the antecubital vein of the arm into vacutainer tubes (BD Bioscience, San Jose, CA, USA) before and after the 8-week intervention. Blood samples were separated by centrifugation at 1,500×g and 4°C for 15 min and stored frozen at -80°C until assayed. Fasting serum glucose level was measured using an automated glucose analyzer (ADVIA 1650, Bayer). Serum insulin (Mercodia, Winston Salem, NC, USA) and irisin (Phoenix Pharmaceuticals, Inc., Burlingame, CA, USA) were assessed using enzyme-linked immunosorbent assay kits. IGF-1 was determined by chemiluminescence immunoassay using the IGF-1 assay kit (Siemens Corporation, UK).

Isokinetic muscle strength (peak torque)

Participants' maximal muscle strength was evaluated at baseline and after 8-week intervention during a laboratory visit. They were seated on a Human/Norm Testing and Rehabilitation System (Computer Sports Medicine, Inc., MA, USA, 2006). Their chests, hips, and thighs were fixed using a belt to minimize the influence of other muscles. The axis of rotation of the knee joint was aligned with that of a dynamometer. After the dynamometer was fixed to the ankle with a strap, anatomical zero was set, and the movable range was set with extension at 0° and flexion at 110°. The participants were provided with sufficient explanation of the test procedure and asked to repeat the flexion/ex-

tension exercise twice to adapt to the test before the measurement. Peak torque was determined by completion of 1 set of 3 flexion/extension 60%/s. Participants had continuous verbal encouragement from supervisors during all tests, and as measurement criteria we selected peak and average torque at each repetition.

Statistical analysis

For all variables, data were expressed as mean±SD and analyzed using PASW Statistics for Windows software version 19.0 (SPSS, version 19.0 for Windows; SPSS, Inc., Chicago, Illinois, USA). The effects of RT and UA supplementation on body composition, blood parameters and muscle strength were tested using two-way ANOVA with repeated measures (2 [groups]×2 [time points]). When a significant group-by-time interaction was detected, within-group differences between mean values were analyzed using a paired t-test. The level of significance was considered p<0.05.

RESULTS

The characteristics of the study participants are shown in Table 1. At baseline, no significant differences in age, height, weight, body mass index, body percentage, or lean body mass were observed between the RT and RT+UA groups. After the RT+UA intervention, body fat percentage was significantly decreased compared to that at baseline in the RT group (p<0.001). Although lean body mass demonstrated a slight increase compared with baseline levels (p>0.05), no significant changes were observed in any characteristics of body composition.

Blood parameters of the study participants are shown in Table 2. At baseline, no significant differences in blood parameters were observed between the RT and RT+UA groups. After RT+UA intervention, IGF-1 and irisin levels were significantly increased compared to baseline in the RT group (p<0.05), while glucose and insulin did not change in either group.

Isokinetic maximal muscle strength of the study partic-

Table 1. Characteristics of participants before and after study interventions

| | RT (n=7) | | RT+UA (n=9) | |
|------------------------|-------------|------------|-------------|-------------------------|
| | Baseline | After 8 wk | Baseline | After 8 wk |
| Age, years | 28.71±5.76 | - | 29.88±4.91 | - |
| Height, cm | 172.57±4.89 | - | 174.09±5.13 | - |
| Weight, kg | 83.15±6.71 | 80.91±6.90 | 80.30±6.00 | 79.89±6.50 |
| BMI, kg/m ² | 27.97±2.58 | 27.20±2.58 | 26.48±1.63 | 26.36±2.03 |
| Body fat, % | 11.95±3.13 | 12.78±3.36 | 11.70±1.80 | 8.70±2.51* [†] |
| LBM, kg | 68.55±6.23 | 68.22±7.03 | 67.75±4.31 | 68.92±3.94 |

Values are expressed as mean±standard deviation. BMI, body mass index; LBM, lean body mass. Within-group differences were analyzed by paired t-test (*p<0.001). Analysis of variance was used for group-by-time interactions ([†]p<0.001).

Table 2. Blood parameters of participants before and after study interventions

| | RT (n=7) | | RT+UA (n=9) | |
|----------------|----------------|----------------|----------------|------------------------------|
| | Baseline | After 8 wk | Baseline | After 8 wk |
| Glucose, mg/dl | 90.00±15.81 | 95.57±15.08 | 84.55±32.11 | 98.11±17.35 |
| Insulin, mg/dl | 18.57±13.66 | 12.97±8.27 | 16.79±6.84 | 13.70±11.78 |
| IGF-1, ng/mL | 179.65±51.47 | 182.54±48.08 | 180.43±59.88 | 221.04±58.29* [†] |
| Irisin, ng/mL | 1461.70±219.88 | 1384.94±159.49 | 1436.49±362.79 | 1613.05±295.25* [†] |

Values are expressed as mean±standard deviation. IGF-I, insulin-like growth factor-I. Within-group differences were analyzed by paired t-test (*p<0.05). Analysis of variance was used for group-by-time interactions ([†]p<0.05).

Table 3. Maximal muscle strength of participants before and after study interventions

| | RT (n=7) | | RT+UA (n=9) | |
|----------|--------------|--------------|--------------|------------------------------|
| | Baseline | After 8 wk | Baseline | After 8 wk |
| MRE, N/m | 266.42±57.13 | 256.28±46.99 | 265.44±42.33 | 281.00±41.07** ^{††} |
| MLE, N/m | 238.28±56.69 | 231.57±42.80 | 237.11±45.97 | 256.44±50.61** [†] |
| MRF, N/m | 166.00±33.02 | 154.14±21.25 | 164.88±20.44 | 179.33±11.50* [†] |
| MLF, N/m | 146.28±20.77 | 139.57±21.25 | 143.55±17.40 | 161.00±14.95*** [†] |

Values are expressed as mean±standard deviation. MRE, maximal right extension; MLE, maximal left extension; MRF, maximal right flexion; MLF, maximal left flexion. Within-group differences were analyzed by paired t-test (*p<0.05, **p<0.01, ***p<0.001). Analysis of variance was used for group-by-time interactions ([†]p<0.05, ^{††}p<0.01).

ipants is shown in Table 3. At baseline, no significant differences in muscle strength were observed between the groups. After RT+UA intervention, maximal right extension ($p < 0.01$), maximal left extension ($p < 0.01$), maximal right flexion ($p < 0.05$), and maximal left flexion ($p < 0.001$) were significantly increased compared to baseline. There were significant group-by-time interactions for maximal right extension ($p < 0.01$), maximal left extension, maximal right flexion, and maximal left flexion ($p < 0.05$) indicating that the increases in isokinetic maximal strength in the RT+UA groups significantly differed from the non-significant changes from baseline in the RT group.

DISCUSSION

In the present study, we demonstrate for the first time that prolonged RT with UA supplementation in men strikingly enhanced physiological changes including a decline in body fat percentage, increase in IGF-1 and irisin levels, and increase in muscle strength. In contrast to, no change in skeletal muscle mass was observed. These findings indicate that RT with UA supplementation may represent a better strategy for increasing skeletal muscle strength via an improvement in plasma IGF-1 and irisin compared to RT alone. Therefore, it appears that UA supplementation can exert a beneficial effect on a combined exercise program, although it may have a mismatched effect on skeletal muscle mass.

Relevant strategies for improving exercise performance include not only positive training methods [28-30], e.g. resistance exercise, but also nutritional protocols [15,31]. Many edible fruits and herbs contain UA, which is a bioactive phyto-constituent [20,32-35] and is also used in anti-obesity herbal medicines [21]. UA appear to strongly inhibit body fat gain in obese humans and animals [26,27]. Similar to these previous results, we observed significant changes in body fat percentage, but no changes in body weight (Table 1). These results suggest that UA supplementation lowered the body fat percentage in men during the study intervention, whereas RT alone did not affect body fat percentage. Therefore, the significant decrease in body fat percentage without change in body weight induced by UA supplementation may be explained by favorable changes in body composition including a decrease in body fat mass [36].

A recent study suggested UA supplementation as a new therapeutic application for skeletal muscle hypertrophy through activation of IGF-1 signaling [6]. In addition, Ogasawara et al. [26] found that UA supplementation stimulated skeletal muscle hypertrophy after RT through Akt-independent activation of mammalian target of rapamycin complex 1 activity, which is known as an important regulator of muscle protein synthesis and hypertrophy [37]. For these reasons, we hypothesized that UA supplementation may improve skeletal muscle hypertrophy via IGF-1 signaling during RT, however, we did not observe substantial gains in lean body mass (Table 1), although a slight, non-significant upward trend was noted. A previous study reported that an increase in lean body mass can stimulate hormones or myokines, which are associated with increased skeletal muscle mass [38]. Additionally, IGF-1 is strongly associated with skeletal muscle mass; is an important hormone involved in skeletal muscle hypertrophy [39]; and regulates cell proliferation, differentiation

[40], and satellite cell function during muscle regeneration [41]. To support our hypothesis that UA exerts effects through increased IGF-1, we observed a significant increase in IGF-1 in the RT+UA group compared to the RT group (Table 2). It appears that UA supplementation may play a role in augmenting IGF-1 without a change in skeletal muscle hypertrophy. Many studies have reported that RT alone [42,43] and UA supplementation [22] significantly impact stimulated IGF-1. These findings indicate that an enhancement of the increase in IGF-1 could be expected in skeletal muscle hypertrophy. However, our training program was not associated with marked skeletal muscle hypertrophy. This discrepancy may be due to our study population, since the men already demonstrated muscle hypertrophy, and thus a further significant increase in muscle mass following 8 weeks of RT alone and RT+UA is not easily achieved. In addition, the adaptation of skeletal muscle hypertrophy, which can lead to the stimulation of other phenomena, may depend on exercise training intensity, frequency, and the duration of physical activity [44]. Therefore, further investigation is needed to consider exercise training protocols and verify the mechanisms underlying UA-associated skeletal muscle hypertrophy in resistance trained-men.

Skeletal muscle has been recently identified as an endocrine organ that can regulate several physiological and metabolic pathways in response to muscle contraction [45]. Several myokines [46] and IGF-1 may be involved in inducing skeletal muscle hypertrophy or suppressing skeletal muscle atrophy [47]. One of these myokines is irisin, which plays a role in energy generation in the skeletal muscle [48]. The exercise-mediated increase in circulating irisin was strongly related with improvement of IGF-1 [6-10]. To clarify the inconsistency in previous findings on irisin, we measured the plasma level of irisin during RT+UA supplementation and found a significant increase in irisin level in the RT+UA group compared to the RT group (Table 2). This finding is the first reported evidence that UA administration leads to increased irisin and that RT alone does not affect this myokine in men. This result raises the possibility that blood levels of IGF-1 and irisin are regulated differently, but the mechanisms underlying their regulation remain unknown.

Finally, we hypothesized that the increase in IGF-1 and irisin would improve maximal muscle strength, since skeletal muscle strength is a quality indicator of skeletal muscle function. UA supplementation during RT significantly enhanced isokinetic muscle strength (Table 3). This result provides another indication that UA may play an important role in muscle strength as well as muscle mass. The finding highlights a novel possibility for obtaining more muscle strength through UA supplementation in men that may perhaps be applicable to other populations who practice RT, such as individuals with muscle atrophy due to aging and/or disease. It is difficult to directly connect increased muscle strength with irisin level following UA supplementation and we cannot exclude the possibility that the increased muscle strength may be related to irisin in combination with IGF-1 or IGF-1 alone. However, these findings suggest that irisin may be involved in exercise-mediated skeletal muscle contraction by increasing the ability to generate maximal muscle force (i.e. muscle strength). The increase in maximal muscle strength in the RT+UA group is highly relevant to health-related fitness benefits, even in well-functioning subjects, since muscle strength helps to

maintain a high level of physical capacity. To our knowledge, no other study has reported independent muscle size and muscle strength responses to UA supplementation in resistance-trained men during RT. The relevance of specific mechanisms underlying these effects of myokines requires further study.

The present study has some limitations. Although significant results were observed in the relatively small number of participants, significant correlations could not be detected in this study. Future studies should aim for a larger number of participants. We provided education about diet to all participants to avoid other diet effects during the study; however, we could not confirm detailed dietary patterns during the study period. Finally, our study was performed in healthy men. In future investigations, the biological effect of UA intervention alone or combined with RT should be performed in individuals with disease states such as muscle weakness and age-and/or disease-related muscle atrophy to further elucidate the muscle strength-enhancing properties of UA.

In conclusion, our results show that an 8-week RT program with UA supplementation can decrease body fat percentage and increase IGF-1, irisin, and skeletal muscle strength without affecting muscle mass in men. Thus, this approach appears to be a promising strategy to improve skeletal muscle strength in men. However, the underlying molecular mechanisms involved in the increase of IGF-1, irisin, and muscle strength achieved through UA supplementation require further investigation.

ACKNOWLEDGEMENTS

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF), and funding was granted by the Ministry of Science, ICT & Future Planning of Korea (R13-2007-023-00000-0, 2011-0028925 and 2012R1A2A1A030-07595), by the Ministry of Education, Science and Technology Korea (2010-0020224) and a 2005 Inje University research grant.

REFERENCES

1. **Yaspelkis BB 3rd.** Resistance training improves insulin signaling and action in skeletal muscle. *Exerc Sport Sci Rev.* 2006; 34:42-46.
2. **Febbraio MA, Pedersen BK.** Contraction-induced myokine production and release: is skeletal muscle an endocrine organ? *Exerc Sport Sci Rev.* 2005;33:114-119.
3. **Mormeneo E, Jimenez-Mallebrera C, Palomer X, De Nigris V, Vázquez-Carrera M, Orozco A, Nascimento A, Colomer J, Lerín C, Gómez-Foix AM.** PGC-1 α induces mitochondrial and myokine transcriptional programs and lipid droplet and glycogen accumulation in cultured human skeletal muscle cells. *PLoS One.* 2012;7:e29985.
4. **Keipert S, Ost M, Johann K, Imber F, Jastroch M, van Schothorst EM, Keijer J, Klaus S.** Skeletal muscle mitochondrial uncoupling drives endocrine cross-talk through the induction of FGF21 as a myokine. *Am J Physiol Endocrinol Metab.* 2014;306:E469-482.
5. **Boström P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Boström EA, Choi JH, Long JZ, Kajimura S, Zingaretti MC, Vind BF, Tu H, Cinti S, Höglund K, Gygi SP, Spiegelman BM.** A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature.* 2012;481:463-468.
6. **Kunkel SD, Elmore CJ, Bongers KS, Ebert SM, Fox DK, Dyle MC, Bullard SA, Adams CM.** Ursolic acid increases skeletal muscle and brown fat and decreases diet-induced obesity, glucose intolerance and fatty liver disease. *PLoS One.* 2012; 7:e39332.
7. **Novelle MG, Contreras C, Romero-Picó A, López M, Diéguez C.** Irisin, two years later. *Int J Endocrinol.* 2013;2013:746281.
8. **Park KH, Zaichenko L, Brinkoetter M, Thakkar B, Sahin-Efe A, Joung KE, Tsoukas MA, Geladari EV, Huh JY, Dincer F, Davis CR, Crowell JA, Mantzoros CS.** Circulating irisin in relation to insulin resistance and the metabolic syndrome. *J Clin Endocrinol Metab.* 2013;98:4899-4907.
9. **Pekkala S, Wiklund PK, Hulmi JJ, Ahtiainen JP, Horttanainen M, Pöglänen E, Mäkelä KA, Kainulainen H, Häkkinen K, Nyman K, Alén M, Herzig KH, Cheng S.** Are skeletal muscle FNDC5 gene expression and irisin release regulated by exercise and related to health? *J Physiol.* 2013;591:5393-5400.
10. **Huh JY, Dincer F, Mesfum E, Mantzoros CS.** Irisin stimulates muscle growth-related genes and regulates adipocyte differentiation and metabolism in humans. *Int J Obes (Lond).* 2014. [Epub ahead of print]
11. **Boström PA, Fernández-Real JM.** Metabolism: Irisin, the metabolic syndrome and follistatin in humans. *Nat Rev Endocrinol.* 2014;10:11-12.
12. **Norheim F, Langleite TM, Hjorth M, Holen T, Kielland A, Stadheim HK, Gulseth HL, Birkeland KI, Jensen J, Drevon CA.** The effects of acute and chronic exercise on PGC-1 α , irisin and browning of subcutaneous adipose tissue in humans. *FEBS J.* 2014;281:739-749.
13. **Moraes C, Leal VO, Marinho SM, Barroso SG, Rocha GS, Boaventura GT, Mafra D.** Resistance exercise training does not affect plasma irisin levels of hemodialysis patients. *Horm Metab Res.* 2013;45:900-904.
14. **Erickson HP.** Irisin and FNDC5 in retrospect: An exercise hormone or a transmembrane receptor? *Adipocyte.* 2013;2:289-293.
15. **Karimian J, Esfahani PS.** Supplement consumption in body builder athletes. *J Res Med Sci.* 2011;16:1347-1353.
16. **Wilson JM, Joy JM, Lowery RP, Roberts MD, Lockwood CM, Manninen AH, Fuller JC, De Souza EO, Baier SM, Wilson SM, Rathmacher JA.** Effects of oral adenosine-5'-triphosphate supplementation on athletic performance, skeletal muscle hypertrophy and recovery in resistance-trained men. *Nutr Metab (Lond).* 2013;10:57.
17. **Yao LH, Meng W, Song RF, Xiong QP, Sun W, Luo ZQ, Yan WW, Li YP, Li XP, Li HH, Xiao P.** Modulation effects of cordycepin on the skeletal muscle contraction of toad gastrocnemius muscle. *Eur J Pharmacol.* 2014;726:9-15.
18. **Kim K, Kim YH, Lee SH, Jeon MJ, Park SY, Doh KO.** Effect of exercise intensity on unfolded protein response in skeletal muscle of rat. *Korean J Physiol Pharmacol.* 2014;18:211-216.
19. **Seo DY, Kwak HB, Lee SR, Cho YS, Song IS, Kim N, Bang HS, Rhee BD, Ko KS, Park BJ, Han J.** Effects of aged garlic extract and endurance exercise on skeletal muscle FNDC-5 and circulating irisin in high-fat-diet rat models. *Nutr Res Pract.* 2014;8:177-182.
20. **Liu J.** Pharmacology of oleanolic acid and ursolic acid. *J Ethnopharmacol.* 1995;49:57-68.
21. **Rao VS, de Melo CL, Queiroz MG, Lemos TL, Menezes DB, Melo TS, Santos FA.** Ursolic acid, a pentacyclic triterpene from *Sambucus australis*, prevents abdominal adiposity in mice fed a high-fat diet. *J Med Food.* 2011;14:1375-1382.
22. **Kunkel SD, Suneja M, Ebert SM, Bongers KS, Fox DK, Malmberg SE, Alipour F, Shields RK, Adams CM.** mRNA expression signatures of human skeletal muscle atrophy identify a natural compound that increases muscle mass. *Cell Metab.* 2011;13:627-638.
23. **Gai L, Cai N, Wang L, Xu X, Kong X.** Ursolic acid induces apoptosis via Akt/NF- κ B signaling suppression in T24 human bladder cancer cells. *Mol Med Rep.* 2013;7:1673-1677.
24. **Chen J, Ko KM.** Ursolic-Acid-Enriched Herba Cynomorii

- Extract Protects against Oxidant Injury in H9c2 Cells and Rat Myocardium by Increasing Mitochondrial ATP Generation Capacity and Enhancing Cellular Glutathione Redox Cycling, Possibly through Mitochondrial Uncoupling. *Evid Based Complement Alternat Med.* 2013;2013:924128.
25. Kim SH, Hong JH, Lee YC. Ursolic acid, a potential PPAR γ agonist, suppresses ovalbumin-induced airway inflammation and Penh by down-regulating IL-5, IL-13, and IL-17 in a mouse model of allergic asthma. *Eur J Pharmacol.* 2013;701:131-143.
 26. Ogasawara R, Sato K, Higashida K, Nakazato K, Fujita S. Ursolic acid stimulates mTORC1 signaling after resistance exercise in rat skeletal muscle. *Am J Physiol Endocrinol Metab.* 2013;305:E760-765.
 27. Souza DR, Gomides RS, Costa LA, Queiroz AC, Barros S, Ortega KC, Mion D Jr, Tinucci T, Forjaz CL. Amlodipine reduces blood pressure during dynamic resistance exercise in hypertensive patients. *Scand J Med Sci Sports.* 2013. [Epub ahead of print]
 28. Seo DY, Lee SR, Kim N, Ko KS, Rhee BD, Han J. Humanized animal exercise model for clinical implication. *Pflugers Arch.* 2014;466:1673-1687.
 29. Seo DY, Lee S, Figueroa A, Kim HK, Baek YH, Kwak YS, Kim N, Choi TH, Rhee BD, Ko KS, Park BJ, Park SY, Han J. Yoga training improves metabolic parameters in obese boys. *Korean J Physiol Pharmacol.* 2012;16:175-180.
 30. Figueroa A, Park SY, Seo DY, Sanchez-Gonzalez MA, Baek YH. Combined resistance and endurance exercise training improves arterial stiffness, blood pressure, and muscle strength in postmenopausal women. *Menopause.* 2011;18:980-984.
 31. de Silva A, Samarasinghe Y, Senanayake D, Lanerolle P. Dietary supplement intake in national-level Sri Lankan athletes. *Int J Sport Nutr Exerc Metab.* 2010;20:15-20.
 32. Verano J, González-Trujano ME, Déciga-Campos M, Ventura-Martínez R, Pellicer F. Ursolic acid from *Agastache mexicana* aerial parts produces antinociceptive activity involving TRPV1 receptors, cGMP and a serotonergic synergism. *Pharmacol Biochem Behav.* 2013;110:255-264.
 33. Kim HY, Choi HR, Lee YJ, Cui HZ, Jin SN, Cho KW, Kang DG, Lee HS. Accentuation of ursolic acid on muscarinic receptor-induced ANP secretion in beating rabbit atria. *Life Sci.* 2014;94:145-150.
 34. Seo DY, Lee S, Figueroa A, Kwak YS, Kim N, Rhee BD, Ko KS, Bang HS, Baek YH, Han J. Aged garlic extract enhances exercise-mediated improvement of metabolic parameters in high fat diet-induced obese rats. *Nutr Res Pract.* 2012;6:513-519.
 35. Seo DY, Lee SR, Kim HK, Baek YH, Kwak YS, Ko TH, Kim N, Rhee BD, Ko KS, Park BJ, Han J. Independent beneficial effects of aged garlic extract intake with regular exercise on cardiovascular risk in postmenopausal women. *Nutr Res Pract.* 2012;6:226-231.
 36. Kazmi I, Afzal M, Rahman S, Iqbal M, Imam F, Anwar F. Antiobesity potential of ursolic acid stearoyl glucoside by inhibiting pancreatic lipase. *Eur J Pharmacol.* 2013;709:28-36.
 37. Apró W, Wang L, Pontén M, Blomstrand E, Sahlin K. Resistance exercise induced mTORC1 signaling is not impaired by subsequent endurance exercise in human skeletal muscle. *Am J Physiol Endocrinol Metab.* 2013;305:E22-32.
 38. Ramakrishnan SN, Lau P, Burke LJ, Muscat GE. Rev-erb β regulates the expression of genes involved in lipid absorption in skeletal muscle cells: evidence for cross-talk between orphan nuclear receptors and myokines. *J Biol Chem.* 2005;280:8651-8659.
 39. Li M, Li C, Parkhouse WS. Age-related differences in the des IGF-I-mediated activation of Akt-1 and p70 S6K in mouse skeletal muscle. *Mech Ageing Dev.* 2003;124:771-778.
 40. Schiaffino S, Mammucari C. Regulation of skeletal muscle growth by the IGF1-Akt/PKB pathway: insights from genetic models. *Skelet Muscle.* 2011;1:4.
 41. Jang YC, Sinha M, Cerletti M, Dall'Osso C, Wagers AJ. Skeletal muscle stem cells: effects of aging and metabolism on muscle regenerative function. *Cold Spring Harb Symp Quant Biol.* 2011;76:101-111.
 42. Velloso CP, Aperghis M, Godfrey R, Blazevich AJ, Bartlett C, Cowan D, Holt RI, Bouloux P, Harridge SD, Goldspink G. The effects of two weeks of recombinant growth hormone administration on the response of IGF-I and N-terminal pro-peptide of collagen type III (P-III-NP) during a single bout of high resistance exercise in resistance trained young men. *Growth Horm IGF Res.* 2013;23:76-80.
 43. Gregory SM, Spiering BA, Alemany JA, Tuckow AP, Rarick KR, Staab JS, Hatfield DL, Kraemer WJ, Maresh CM, Nindl BC. Exercise-induced insulin-like growth factor I system concentrations after training in women. *Med Sci Sports Exerc.* 2013;45:420-428.
 44. Schoenfeld BJ. Is there a minimum intensity threshold for resistance training-induced hypertrophic adaptations? *Sports Med.* 2013;43:1279-1288.
 45. Keipert S, Ost M, Johann K, Imber F, Jastroch M, van Schothorst EM, Keijer J, Klaus S. Skeletal muscle mitochondrial uncoupling drives endocrine cross-talk through the induction of FGF21 as a myokine. *Am J Physiol Endocrinol Metab.* 2014;306:E469-482.
 46. Pedersen BK, Akerström TC, Nielsen AR, Fischer CP. Role of myokines in exercise and metabolism. *J Appl Physiol (1985).* 2007;103:1093-1098.
 47. Philippou A, Halapas A, Maridaki M, Koutsilieris M. Type I insulin-like growth factor receptor signaling in skeletal muscle regeneration and hypertrophy. *J Musculoskelet Neuronal Interact.* 2007;7:208-218.
 48. Villarroya F. Irisin, turning up the heat. *Cell Metab.* 2012;15:277-278.