

Original Article

## Sodium butyrate has context-dependent actions on dipeptidyl peptidase-4 and other metabolic parameters

Eun-Sol Lee<sup>1,#</sup>, Dong-Sung Lee<sup>2,#</sup>, Prakash Raj Pandeya<sup>1</sup>, Youn-Chul Kim<sup>1</sup>, Dae-Gil Kang<sup>3,4</sup>, Ho-Sub Lee<sup>3,4</sup>,  
Byung-Chul Oh<sup>5,6</sup>, and Dae Ho Lee<sup>7,8,\*</sup>

<sup>1</sup>College of Pharmacy, Wonkwang University, Iksan 54538, <sup>2</sup>College of Pharmacy, Chosun University, Gwangju 61452, <sup>3</sup>Hanbang Body Fluid Research Center, Wonkwang University, Iksan 54538, <sup>4</sup>College of Oriental Medicine and Professional Graduate School of Oriental Medicine, Wonkwang University, Iksan 54538, <sup>5</sup>Lee Gil Ya Cancer and Diabetes Institute, College of Medicine, Gachon University, Incheon 21999, <sup>6</sup>Department of physiology, College of Medicine, Gachon University, Incheon 21999, <sup>7</sup>Department of Internal Medicine, Gachon University Gil Medical Center, Incheon 21565, <sup>8</sup>Department of Internal Medicine, Gachon University College of Medicine, Incheon 21999, Korea

### ARTICLE INFO

Received March 13, 2017  
Revised May 29, 2017  
Accepted June 20, 2017

### \*Correspondence

Dae Ho Lee  
E-mail: drhormone@naver.com

### Key Words

Dipeptidyl peptidase 4  
HepG2 cells  
Mesangial cells  
Obesity  
Sodium butyrate

#These authors contributed equally to this work.

**ABSTRACT** Sodium butyrate (SB) has various metabolic actions. However, its effect on dipeptidyl peptidase 4 (DPP-4) needs to be studied further. We aimed to evaluate the metabolic actions of SB, considering its physiologically relevant concentration. We evaluated the effect of SB on regulation of DPP-4 and its other metabolic actions, both *in vitro* (HepG2 cells and mouse mesangial cells) and *in vivo* (high fat diet [HFD]-induced obese mice). Ten-week HFD-induced obese C57BL/6J mice were subjected to SB treatment by adding SB to HFD which was maintained for an additional 16 weeks. In HepG2 cells, SB suppressed DPP-4 activity and expression at sub-molar concentrations, whereas it increased DPP-4 activity at a concentration of 1,000  $\mu$ M. In HFD-induced obese mice, SB decreased blood glucose, serum levels of insulin and IL-1 $\beta$ , and DPP-4 activity, and suppressed the increase in body weight. On the contrary, various tissues including liver, kidney, and peripheral blood cells showed variable responses of DPP-4 to SB. Especially in the kidney, although DPP-4 activity was decreased by SB in HFD-induced obese mice, it caused an increase in mRNA expression of TNF- $\alpha$ , IL-6, and IL-1 $\beta$ . The pro-inflammatory actions of SB in the kidney of HFD-induced obese mice were recapitulated by cultured mesangial cell experiments, in which SB stimulated the secretion of several cytokines from cells. Our results showed that SB has differential actions according to its treatment dose and the type of cells and tissues. Thus, further studies are required to evaluate its therapeutic relevance in metabolic diseases including diabetes and obesity.

## INTRODUCTION

Dipeptidyl peptidase 4 (DPP-4) is widely expressed on the surface of many cell types and is involved in the dipeptidyl enzymatic metabolism of incretin hormones [1]. The dipeptidyl peptidase enzyme activity is blocked by DPP-4 inhibitors for the treatment of type 2 diabetes mellitus (T2DM) [2]. However, the factors and mechanisms that control DPP-4 activity are very complex and need to be clarified further; hence, further studies are required

even after ten years of widespread use of DPP-4 inhibitors in diabetes clinics [3]. Further, as it is associated with insulin resistance and diabetes, DPP-4 expression is increased in adipose tissue and its stromal vascular dendritic cells/macrophages [4] and peripheral blood T cells [5]. Some studies have shown that glucose level affects DPP-4 activity and expression per se [6]. We have also reported that T cell DPP-4 expression, serum soluble DPP-4 and DPP-4 activity were increased in patients with T2DM, and the increase was related to glycemic status [5]. Also, DPP-4 can be



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.  
Copyright © Korean J Physiol Pharmacol, pISSN 1226-4512, eISSN 2093-3827

**Author contributions:** E.S.L., D.S.L., P.R.P., Y.C.K., and D.H.L. designed the study; participated in the acquisition, analysis, and interpretation of the data; and contributed to writing the manuscript. Y.C.K., B.C.O., D.G.K., and H.S.L. participated in the acquisition, analysis, and interpretation of the data; and contributed to the writing of the manuscript. All authors edited the manuscript and approved the final version.

regulated in a cell- or tissue-specific manner. In Caco2 cells, high glucose concentrations suppress DPP-4 gene expression, resulting in decreased DPP-4 enzymatic activity [7]. In HepG2 cells, DPP-4 activity and gene expression were found to be increased by high dose of sodium butyrate (SB) [8].

Recently, butyric acid and its solute form SB have been gaining increasing attention due to the metabolic actions of short-chain fatty acids (SCFAs), which are naturally produced in the distal small intestine and colon through fermentation of resistant starch, dietary fiber, and other low-digestible polysaccharides [9]. SB and other SCFAs, suppress histone deacetylases (HDACs), thereby regulating cell growth, differentiation and apoptosis in multiple cells [10]. In dietary obese C57BL/6J mice, SB was shown to suppress weight gain independent of food intake suppression. Also, except for the intestinal lumen, most of SCFAs in tissue and blood exist in their submolar concentrations [11], which is in contrast to the high molar concentration of SB that was used in *in vitro* experiments. Many studies have suggested that butyrate has pleiotropic effects through the modification of chromatin plasticity, histone acetylation, histone phosphorylation and/or phosphorylation of high-mobility group proteins [8]. However, some studies have shown that SB has dual effects on cell growth, cell cycle distribution, and gene expression in hepatocellular carcinoma cells depending on the concentration [12] and cell-specific actions depending on the cell type [13], which warrant further studies regarding the previously reported effect of SB on DPP-4. To understand clinical implications of the pleiotropic actions of butyrate, it seems necessary to address the context-dependent actions of butyrate. In this study, we evaluated the effect of SB on regulation of DPP-4 activity and expression in cells and high fat diet (HFD)-induced obese mice to better characterize the metabolic actions of SB.

## METHODS

### Materials

RPMI 1640, fetal bovine serum (FBS) and other cell culture reagents were purchased from Gibco BRL Co. (Grand Island, NY, USA). All primary and secondary antibodies, unless otherwise stated, were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Anti-acetyl-H3 and anti-acetyl-H4 antibodies were purchased from Millipore (Bedford, MA, USA). The enzyme-linked immunosorbent assay (ELISA) kits for interleukin (IL)-1 $\beta$ , IL-6, and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) were purchased from R&D Systems (Minneapolis, MN, USA). Histone extraction kit was obtained from Abcam (Cambridge, UK). Other chemicals, unless stated otherwise, were obtained from Sigma Chemical Co. (St. Louis, MO, USA).

### Cell culture

HepG2 and mesangial cells were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA). HepG2 cells were maintained in RPMI 1640 supplemented with 10% heat-inactivated fetal bovine serum (FBS), penicillin G (100 units/mL), streptomycin (100 mg/mL), and L-glutamine (2 mmol/L), at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. Mouse mesangial cells (ATCC, Manassas, VA, USA) were maintained in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (DMEM/F12) supplemented with 10% heat-inactivated fetal bovine serum (FBS), with other culture conditions being similar to HepG2 cells. Most experiments were performed in mesangial cells after 3 days of incubation with SB, unless stated otherwise.

### Animal experiments

The animal procedures were conducted in accordance with the 1985 (revised 1996) Guidelines for the Care and Use of Laboratory Animals of the U.S. National Institutes of Health and were approved by the Institutional Animal Care and Utilization Committee for Medical Science of Wonkwang University (Approval NO. WKU15-44). C57BL/6J mice (n $\geq$ 9 in each group) were obtained from Orient Bio (Sungnam, Kyungkido, South Korea). High-fat diet (HFD; 60% of total calories) was purchased from Research Diets, Inc (New Brunswick, NJ, USA, Cat. D12492). All animals were bred and housed in standard cages in a climate-controlled room at an ambient temperature of 23 $\pm$ 2°C. C57BL/6J mice were fed normal chow diet (NCD) or HFD for 10 weeks. Then, HFD-induced obese mice were divided into two groups: HFD only or HFD+SB. In the "HFD+SB" group, mice were kept on HFD with SB incorporated at 5% wt/wt (n=9), while in the "HFD only" group (HFD, n=9) and the "control" group (Control, n=9), mice were kept on the same HFD or NCD, respectively. At 16 weeks thereafter, mice from the three experimental groups were sacrificed. Blood and tissues including liver and kidney were collected appropriately. To prepare plasma for the measurement of active GLP-1, we collected blood in ice-cooled ethylenediaminetetraacetic acid (EDTA)-plasma tubes which contained a DPP-4 inhibitor (10  $\mu$ l per milliliter of blood) obtained from EMD Millipore Corporation (St. Charles, Missouri, USA) and we mixed the blood gently with the reagents. The tubes were maintained in an ice bath and centrifuged within one hour at 1000 x g for 10 minutes in a refrigerated centrifuge. Specimens were stored at -70 °C until assayed.

### Western blot analysis

Cells or mouse tissues were washed with PBS and lysed using RIPA lysis buffer containing 25 mmol/L Tris-HCl buffer (pH 7.6), 150 mmol/L NaCl, 1% NP-40, 1% sodium deoxycholate, and 0.1% SDS. The protein concentration was determined using Bradford

Assay Reagent (Bio-Rad, PA, USA). An equal amount of protein from each sample was resolved using sodium dodecyl sulfate-polyacrylamide gel electrophoresis and it was then electrophoretically transferred onto a Hybond-enhanced chemiluminescence nitrocellulose membrane (Bio-Rad, PA, USA). The membrane was blocked with 5% skim milk and incubated with primary antibodies at 4°C overnight. The bands were visualized with enhanced chemiluminescence and quantified by densitometry. Representative blots from three independent experiments are presented in the figures. Nuclear and cytoplasmic extracts of cells were prepared using NE-PER nuclear and cytoplasmic extraction reagents (Pierce Biotechnology, Rockford, IL USA), respectively.

### DPP-4 activity assay

DPP-4 activity was determined according to the cleavage rate of p-nitroaniline (pNA; Sigma) using the synthetic substrate H-glycyl-prolyl-p-nitroanilide (Gly-Pro-pNA; Sigma) [14]. Briefly, 100 µl of assay mixture was prepared containing 50 mM Tris (pH 8.0) and the protein sample from cells or mice tissues, or cell culture medium. The reaction was initiated by the addition of a 5 mM Gly-Pro-pNA substrate solution to a final concentration of 2.5 mM. After incubation for 10~30 min at 37°C, the absorbance of the sample in each well was measured at 405 nm using a plate reader. DPP-4 activity was expressed as the amount of cleaved pNA released per minute per ml (nmol/min/ml) for the culture medium and as the amount of pNA released per minute per weight (nmol/min/mg) for cells/tissues.

### Real-time PCR

RNA was extracted from cells or tissues using an RNeasy Mini Kit (Qiagen). For real-time polymerase chain reaction (PCR), first-strand complementary DNA (cDNA) was synthesized from 1 µg total RNA using an Advantage RT-for-PCR Kit (Takara Korea Biomedical Inc., Seoul, Korea). Relative messenger RNA levels were determined by real-time PCR using a Brilliant II SYBR Green QPCR Master Mix Kit (Stratagene) and an Mx3000P thermal cycler (Stratagene). All cDNA levels were normalized to the level of glyceraldehyde 3-phosphate dehydrogenase (GAPDH). The primer sequences were designed using PrimerQuest (Integrated DNA Technologies, Cambridge, MA, USA).

### Cytokine and monocyte chemotactic protein (MCP)-1 measurements

The serum levels of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and MCP-1 in each sample were determined using a commercially available kit from R&D Systems (Minneapolis, MN, USA). The assay was performed according to the manufacturer's instructions.

### Histone extraction

Histone was extracted from cells using a commercially available Histone extraction kit from Abcam (Cambridge, UK). The assay was performed according to the manufacturer's instructions.


### Statistical analysis

The data were expressed as the mean  $\pm$  standard error (S.E.) of at least three independent experiments or at least five independent samples and for at least six mice in each group. To compare three or more groups, one-way analysis of variance (ANOVA) followed by the Newman-Keuls *post hoc* test was used. Statistical analysis was performed using GraphPad Prism software, version 3.03 (GraphPad Software Inc., San Diego, CA, USA).

## RESULTS

### SB has dual effects on regulation of DPP-4 in HepG2 cells

To evaluate the effect of SCFAs on DPP-4 activity, we incubated HepG2 cells with SB, sodium acetate (SA), or sodium propionate (SP) for 3 days and measured DPP-4 activity in cell lysates. SB inhibited DPP-4 activity at the concentration range of 1 to 100 µM, whereas the DPP-4 activity was increased when the cells were treated with high concentration (1,000 µM) of SB, at which HDAC activity was reported to be inhibited significantly [12]. SP treatment of HepG2 cells showed similar but weaker effects on DPP-4 activity compared with SB. However, SA increased DPP-4 activity in HepG2 cells at 1 and 10 µM concentration. After we observed prominent and reciprocal effects of SB on DPP-4 activity in HepG2 cells, we focused on the effect of SB and evaluated the effects of SB on mRNA and protein expressions of DPP-4 and the acetylation of histone proteins in HepG2 cells. The expressions of both DPP-4 protein and mRNA were significantly increased in HepG2 cells treated with high concentration of SB, whereas both DPP-4 protein and mRNA expressions were down-regulated in HepG2 cells treated with a lower concentration of SB (100 µM) (Figs. 1A-1E). We also investigated the expression of acetylated histone 3 (Acetyl-H3) and acetylated histone 4 (Acetyl-H4), as surrogate markers of HDAC inhibition, and we found the expression levels of Acetyl-H3 and Acetyl-H4 were consistently increased in HepG2 cells treated with high concentration of SB, without suppression at a lower concentration of SB (Fig. 1F). These results suggest that the suppression of DPP-4 activity and expression in HepG2 cells by relatively lower concentrations of SB is not related to the inhibitory action of SB on HDAC.



**Fig. 1. The effects of SB on DPP-4 regulation (A~E) or acetyl-H3 and acetyl-H4 acetylation (F) in HepG2 cells.** HepG2 cells were treated with SB (0, 1, 10, 100, 1,000 μM) for 72 h. DPP-4 activity was determined as described in the “Methods” section. Histone was extracted using a histone extraction kit. Data represent means±S.E. \* $p < 0.05$ , \*\* $p < 0.01$ , and \*\*\* $p < 0.001$  vs. untreated cells.


### SB does not suppress DPP-4 activity in HepG2 cells under high glucose condition

Our previous study showed that serum DPP-4 activity and soluble DPP-4 level were increased in patients with T2DM [5]. These previous findings prompted us to evaluate whether the exposure of HepG2 cells to high glucose increases DPP-4 activity or not. DPP-4 activity and mRNA expression were markedly increased when the cells were incubated for 6 or more days under high glucose (30 mM) condition compared with that in HepG2 cells incubated in control media containing 11 mM glucose (Figs. 2A-2C). To explore if SB has a suppressive effect on the high glucose-induced increase in DPP-4 activity, we incubated HepG2 cells in high glucose media supplemented with SB. Interestingly, low concentration of SB did not suppress DPP-4 activity under high glucose condition. However, contrary to the findings in control media, SB at the low concentration range increased DPP-4 activity in HepG2 cells under high glucose conditions within 2~6 days (Figs. 2D-2F). The results suggest that SB, at its physiologically relevant concentration, has a context-dependent effect on DPP-4, at least in HepG2 cells.


### SB suppresses DPP-4 activity and improves metabolic abnormalities and inflammatory cytokine profiles in HFD-induced obese mice

Before studying the *in vivo* effect of SB on HFD-induced obese mice, we checked the serum DPP-4 activity in C57BL/6J mice that were fed HFD for different periods from 0 to 24 weeks. Each set of mice ( $n=6$  each) fed HFD for the indicated time period, as shown in Fig. 3A, was matched for age at sacrifice to exclude the effect of ageing on DPP-4 activity. Serum DPP-4 activity was increased significantly from 6 weeks of HFD in the experimental mice, and thereafter continuously across the period of HFD feeding (Fig. 3A). Then, to evaluate the metabolic effect of SB on DPP-4, and in obesity and insulin resistance, we administered SB to HFD-induced obese mice that had been on a HFD for 10 weeks. The HFD-induced obese group showed an additional continuous increase in body weight (BW) during the extended HFD period of 16 weeks, while SB treatment in HFD-fed mice (HFD+SB) suppressed the increase in BW (Fig. 3B). In addition, both blood glucose and serum insulin levels were lower after 16 weeks of SB treatment compared with those in non-treated HFD-induced obese mice (Figs. 3C, 3D). The dietary intake was not influenced by SB treatment (data not shown).

Additionally, we evaluated the effects of SB on serum DPP-4 activity and mRNA expression levels of DPP-4, TNF- $\alpha$ , IL-



**Fig. 2. The effects of SB on DPP-4 regulation in HepG2 cells and culture medium under co-stimulation with normal or high glucose condition (A~F).** HepG2 cells were treated with glucose (11 and 30 mM) for up to 8 days (A~C). HepG2 cells were treated with glucose (11 and 30 mM) for 2 days (D), 4 days (E), and 6 days (F). DPP-4 activity was determined as described in the “Methods” section. Data represent means±S.E. \*p<0.05, \*\*p<0.01 \*\*\*p<0.001 vs. glucose 11 mM treated cells.



**Fig. 3. The effects of SB on DPP-4 activity (A), body weight (B), blood glucose (C), and insulin secretion (D) in HFD-induced obese mice.** C57BL/6J mice were fed NCD (Control, n=9) or 60% high-fat diet (HFD) for 10 weeks. Mice in the control group were fed NCD for the additional 16-week period (Control, n=9). HFD-induced obese mice were divided into two groups; the mice were kept on HFD with SB incorporated at 5% wt/wt (HFD+SB, n=9) or HFD only (HFD, n=9) for the additional 16-week period. Data represent means±S.E. \*p<0.05, \*\*p<0.01, and \*\*\*p<0.001 vs. Control group; #p<0.05, ##p<0.01, and ###p<0.001 vs. HFD group.

IL-1 $\beta$ , IL-6, and MCP-1 in peripheral blood cells (PBCs) from the experimental mice. While DPP-4 mRNA expression in PBCs was not different between the experimental groups, increased serum DPP-4 activity in HFD-induced obese mice was suppressed by SB treatment (Figs. 4A, 4B). In addition, mRNA expressions of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and MCP-1 in PBCs were down-regulated by SB treatment in HFD-fed mice (Figs. 4C-4F). Among the three cytokines that we measured in the serum obtained from the experimental groups (Figs. 4G-4I), only the IL-1 $\beta$  level was significantly increased by HFD feeding, and SB treatment significantly suppressed the increase in serum IL-1 $\beta$  (Fig. 4H).


### SB has different effects on DPP-4 and cytokines in the liver and kidney from HFD-induced obese mice

Among tissues, liver and kidney were reported to have a very high DPP-4 activity [15,16]. Therefore, after 16 weeks of SB treatment in HFD-induced obese mice, we investigated the effects of SB treatment on mRNA expression of DPP-4 and cytokines and


DPP-4 activity in liver (Figs. 5A-5F) and kidney (Figs. 5G-5L) obtained from the three experimental groups. Contrary to serum, DPP-4 activity and mRNA expression in the liver were not affected by SB treatment in HFD-induced obese mice. However, the treatment of HFD-induced obese mice with SB suppressed the increase in hepatic mRNA levels of TNF- $\alpha$  (Fig. 5C) and MCP-1 (Fig. 5F).

In the kidney (Figs. 5G-5L), DPP-4 activity and mRNA expression were increased in the HFD group compared with the control group, and these increases were suppressed in the HFD+SB group (Figs. 5G, 5H). However, to our surprise, the renal mRNA expression of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 was significantly increased by SB treatment in HFD-fed mice (Figs. 5I-5L).

Considering the regulating effect of SB on DPP-4 we measured the plasma active GLP-1 level in the experimental mice; however, plasma active GLP-1 levels were comparable between the groups (data not shown).



**Fig. 4. The effects of SB on DPP-4 regulation (A, B), and TNF- $\alpha$  (C, G), IL1 $\beta$  (D, H), IL-6 (E, I), and MCP-1 (F) production in peripheral blood cells (PBCs) or serum of HFD-induced obese mice (as in Fig. 3).** At 16 weeks after combination with SB in HFD-induced obese mice, the mice were sacrificed and the PBCs were collected (B-F). Data represent means $\pm$ S.E. \* $p$ <0.05, \*\* $p$ <0.01, and \*\*\* $p$ <0.001 vs. Control group; # $p$ <0.05, ## $p$ <0.01, and ### $p$ <0.001 vs. HFD group.




**Fig. 5.** The effects of SB on DPP-4 regulation (A, B, G, H), and TNF- $\alpha$  (C, I), IL-1 $\beta$  (D, J), IL-6 (E, K), and MCP-1 (F, L) production in the liver or kidney tissues of HFD-induced obese mice (as in Fig. 3). Data represent means $\pm$ S.E. \* $p$ <0.05, and \*\* $p$ <0.01 vs. Control group; # $p$ <0.05, ## $p$ <0.01, and ### $p$ <0.001 vs. HFD group.

### SB upregulates DPP-4 and increases cytokine secretion in cultured mouse mesangial cells

Finally, to better characterize different effects of SB depending on the concentration and tissue, an additional experiment was conducted using a mesangial cell line. Mesangial cells were treated with indicated concentrations (0~1,000  $\mu$ M) of SB and incubated for 3 days. First, DPP-4 activity and mRNA level were increased by SB at 100 and 1,000  $\mu$ M (Figs. 6A, 6B). Furthermore, secretion of TNF- $\alpha$ , IL-1 $\beta$ , and MCP-1 from cultured mesangial cells was increased significantly at high concentration of SB, whereas IL-6 production was not affected (Figs. 6C-6F).

### DISCUSSION

DPP-4 is well known for its pivotal role in regulation of glycemia *via* catabolism of incretin peptides, and it plays many other roles in cell-matrix interactions and in the regulation of functional peptides [16,17]. DPP-4 expression has been shown to be influenced by hypoxia, with hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) being a strong inducer of DPP-4 gene and protein [18]. Besides HIF-1 $\alpha$ , hepatocyte nuclear factor-1 $\alpha$  (HNF-1 $\alpha$ ), interferons, retinoic acid, various cytokines, and advanced glycation end products have been shown to activate DPP-4 [19-26]. Therefore, DPP-4 has implications for a potential pathophysiological role not only in metabolic disorders, but also in other disorders, including



**Fig. 6.** The effects of SB on DPP-4 regulation (A, B), and TNF- $\alpha$  (C), IL-1 $\beta$  (D), IL-6 (E), and MCP-1 (F) production in mesangial cells. Mesangial cells were treated with SB (0, 1, 10, 100, 1,000  $\mu$ M) for 72 h. Data represent means $\pm$ S.E. \* $p$ <0.05, \*\* $p$ <0.01, and \*\*\* $p$ <0.001 vs. untreated cells.

inflammatory diseases. Thus, further studies on factors regulating *in vivo* DPP-4 activity in both endogenous and exogenous manners seem necessary.

SB, being also known as an HDAC inhibitor, has been receiving much attention because of its wide range of metabolic actions and anti-inflammatory and anti-cancer effects [14,27-34]. Although a previous study demonstrated that SB at a high concentration largely increases DPP-4 activity in HepG2 cells, it was beyond the physiologic range [8]. Other studies revealed that the responsiveness of different cancer cell lines to SB varied greatly. Especially, Jiang et al. [12] reported that while a low concentration of SB promoted cell proliferation and suppressed p21 Cip1 expression, a high concentration of SB induced growth arrest and cell apoptosis, accompanied by p21 Cip1 up-regulation and poly (ADP-ribose) polymerase cleavage. However, the study showed that HDAC inhibition by SB in cancer cell lines including HepG2 cells was clearly dose-dependent from low to high concentrations of SB, indicating differential effects of SB on between HDAC inhibition and the regulation of cellular growth.

Metabolic actions of SB have recently been revealed intensively [11,28,30,32]; however, it needs to be investigated further with regard to DPP-4. The present study evaluated not only the effect of SB on DPP-4 activity *in vitro* and *in vivo*, but also metabolic factors affecting DPP-4 activity. We found that SB suppressed DPP-4 activity at concentrations of 1–100  $\mu$ M in a concentration-dependent manner, but the DPP-4 activity was increased at a high concentration of SB (1,000  $\mu$ M). It has already been reported that a high level of SB (5 mM) increased DPP-4 activity in HepG2 cells [17]; however, there is no report suggesting that a low level of SB decreases DPP-4 activity. Previous studies revealed that SB has a suppressive effect on HDACs consistently [12,31]. We also observed that SB increased Acetyl-H3 in a dose-dependent manner,

in contrast to its effect on DPP-4. Thus, our study showed that DPP-4 has dual effects on DPP-4 activity in HepG2 cells, but not on HDAC activity. Our results are in line with a previous report, which showed that distinct dual effects of SB on cell growth did not match its inhibitory effect on HDAC in hepatic cancer cell lines [12].

We also evaluated the effect of high glucose on DPP-4 activity in HepG2 cells. Long-term exposure of HepG2 cells under the high glucose condition led to a significant increase in DPP-4 activity in the cells. Interestingly enough, in contrast to the results under a relatively low glucose condition (11 mM), under the high glucose condition, SB treatment of HepG2 cells with SB at low concentration (10–100  $\mu$ M) did not lower, but increased DPP-4 activity, even earlier than high glucose condition only. Thus, SB has different effects on cells in context-dependent manners.

We also evaluated HFD on serum DPP-4 activity in mice. Serum DPP-4 activity was increased continuously from 6 weeks of HFD. This is the first study to show the time course of serum DPP-4 activity in HFD-induced animal model for a long-term period. In our previous study, we showed that serum soluble DPP-4 and DPP-4 activity were increased in patients with T2DM [5], while other researchers reported that lower DPP-4 levels following exercise training plus weight loss were related to increased insulin sensitivity in adults with metabolic syndrome [3,35]. Thus, our results and others indicate that metabolic dysregulation itself lead to an increase in serum DPP-4 activity which is actually the therapeutic target of diabetes for incretin effects.


Butyrate increased mitochondrial respiration at the cellular level, through the regulation of energy metabolism-related genes including peroxisome proliferator-activated receptor (PPAR), PPAR $\gamma$  coactivator-1, and carnitine palmitoyltransferase 1b at the molecular level [11,36,37]. In addition, mice group fed a HFD



including 5% SB showed body weight maintenance, muscle increment and blood glucose decrement [11]. To the best of our knowledge, there is no report about SB with respect to the *in vivo* regulation of DPP-4 activity. Therefore, to address this issue, we applied the same diet condition (60% HFD containing 5% SB) as the study performed by Gao et al. [11], and we obtained similar results, especially with respect to body weight and blood glucose. In addition, serum DPP-4 activity in HFD-induced obese mice was significantly suppressed in the group that received 5% SB mixed in the HFD. These results suggested that SB may be effective in preventing obesity and insulin resistance, with one of mechanisms being the regulation of DPP-4 activity. However, this suggestion was not supported by GLP-1 measurement data, which showed no difference in plasma active GLP-1 levels between the groups in the present study. SCFAs, specifically acetate and propionate, but not butyrate, has been shown to increase GLP-1 secretion, probably *via* a SCFA receptor free fatty acid receptor 2 (FFAR2), in experiments using colonic cell culture system and cecal SCFA infusion to mice [38,39]. However, orally ingested SCFAs are absorbed in the small intestine and do not reach the cecum. Thus, oral administration of SCFAs was not shown to increase plasma GLP-1 level in mice [38]. If butyrate inhibits DPP-4 activity in whole body at a range of its physiologic concentrations, one would expect an elevation of circulating active GLP-1 level in SB-treated mice. However, as we observed, tissue- and context-dependent actions of SB might affect the results in the present study. Further studies are required on this issue.

The liver and kidney have significantly higher DPP-4 activities [3]. In addition, the main source of endogenous DPP-4 has been reported to be micro-vascular endothelial cells of some tissues such as liver and kidney [3,15,16,40]. To investigate the metabolic

regulatory action of SB, we also measured on the regulation of AMP-activated protein kinase (AMPK), peroxisome proliferator-activated receptor gamma coactivator (PGC)1- $\alpha$ , and PGC1 expression in the liver tissues of HFD-induced obese mice. AMPK plays a key role in hepatic lipid metabolism, and it is generally known that high-fat diets reduce AMPK activity in the liver tissue [41,42]. However, interestingly, our previous study showed that the phosphorylation of AMPK increased in the liver tissues of HFD-induced obese mice [43]. In the present study, the phosphorylation of AMPK was also significantly increased in the liver tissue of HFD-induced obese mice. And SB treatment in HFD-fed mice (HFD+SB) was increasingly induced the phosphorylation of AMPK more than HFD-fed mice group (Supplementary Fig. 1A). Furthermore, PGC1- $\alpha$ , and PGC1 expression were increased by SB treatment in the liver tissues of HFD-induced obese mice (Supplementary Fig. 1B). Other previous studies also indicated that PGC1-AMPK signaling contributes to improve the metabolic effects [44,45]. Therefore, these results suggested that the regulation of AMPK, PGC1- $\alpha$ , and PGC1 expression might be shown a major metabolic function of SB. In addition, DPP-4 inhibitor may have renoprotective effects on diabetic kidney disease, regardless of its glucose-lowering effects [46]. In this study, SB treatment of HFD-induced obese mice decreased DPP-4 activity and mRNA expression in the kidney, whereas renal TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 mRNA levels were increased. Therefore, we also checked the effects of SB using a mesangial cell line. In mesangial cells, the dual effects of SB on DPP-4 activity were not observed, but DPP-4 activity was increased by treatment with a high concentration of SB. Furthermore, SB treatment of this mesangial cell line increased DPP-4 mRNA expression and stimulated the secretion of several pro-inflammatory cytokines (TNF- $\alpha$ , IL-1 $\beta$ , and MCP-1)



- Collectively, SB has a regulatory action on DPP-4 with some beneficial metabolic effects in cells and HFD-induced obese mice, while it has also differential metabolic actions according to its treatment dose and the type of cells and tissues.

**Fig. 7.** The pathway scheme of the effects of SB on the regulation of DPP-4 as well as other differential actions in HepG2 cells, mouse mesangial cells, and HFD-induced obese mice in context-dependent manners.

in a dose-dependent manner. Thus, although many studies have shown that SB has metabolically beneficial effects and anti-inflammatory effects [34], we found that SB has quite differentiated effects in the kidney for HFD-induced obese mice and mesangial cells. Hence, further studies on the context-dependent actions of SB are needed.

In conclusion, we demonstrated various effect of SB on the regulation of DPP-4 as well as other differential actions in HepG2 cells, mouse mesangial cells, and HFD-induced obese mice in context-dependent manners (Fig. 7). SB has a regulatory action on DPP-4 with some beneficial metabolic effects in cells and HFD-induced obese mice, while it has also differential metabolic actions according to its treatment dose and the type of cells and tissues. Therefore, further studies are required to evaluate its therapeutic relevance in metabolic diseases including diabetes and obesity and to determine safe and effective dosing of SB treatment for the metabolic diseases and their complications.

## ACKNOWLEDGEMENTS

This study was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Korea (HI14C1135).

## CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

## SUPPLEMENTARY MATERIALS

Supplementary data including one figures can be found with this article online at <http://pdf.medrang.co.kr/paper/pdf/Kjpp/Kjpp021-05-06-s001.pdf>.

## REFERENCES

- Morimoto C, Schlossman SF. The structure and function of CD26 in the T-cell immune response. *Immunol Rev*. 1998;161:55-70.
- Choe EY, Cho Y, Choi Y, Yun Y, Wang HJ, Kwon O, Lee BW, Ahn CW, Cha BS, Lee HC, Kang ES. The effect of DPP-4 inhibitors on metabolic parameters in patients with type 2 diabetes. *Diabetes Metab J*. 2014;38:211-219.
- Kim NH, Yu T, Lee DH. The nonglycemic actions of dipeptidyl peptidase-4 inhibitors. *Biomed Res Int*. 2014;2014:368703.
- Zhong J, Rao X, Deiluiis J, Braunstein Z, Narula V, Hazey J, Mikami D, Needleman B, Satoskar AR, Rajagopalan S. A potential role for dendritic cell/macrophage-expressing DPP4 in obesity-induced visceral inflammation. *Diabetes*. 2013;62:149-157.
- Lee SA, Kim YR, Yang EJ, Kwon EJ, Kim SH, Kang SH, Park DB, Oh BC, Kim J, Heo ST, Koh G, Lee DH. CD26/DPP4 levels in peripheral blood and T cells in patients with type 2 diabetes mellitus. *J Clin Endocrinol Metab*. 2013;98:2553-2561.
- Ryskjaer J, Deacon CF, Carr RD, Krarup T, Madsbad S, Holst J, Vilsbøll T. Plasma dipeptidyl peptidase-IV activity in patients with type-2 diabetes mellitus correlates positively with HbA1c levels, but is not acutely affected by food intake. *Eur J Endocrinol*. 2006;155:485-493.
- Gu N, Tsuda M, Matsunaga T, Adachi T, Yasuda K, Ishihara A, Tsuda K. Glucose regulation of dipeptidyl peptidase IV gene expression is mediated by hepatocyte nuclear factor-1alpha in epithelial intestinal cells. *Clin Exp Pharmacol Physiol*. 2008;35:1433-1439.
- Böhm SK, Gum JR Jr, Erickson RH, Hicks JW, Kim YS. Human dipeptidyl peptidase IV gene promoter: tissue-specific regulation from a TATA-less GC-rich sequence characteristic of a housekeeping gene promoter. *Biochem J*. 1995;311:835-843.
- Freeland KR, Wolever TM. Acute effects of intravenous and rectal acetate on glucagon-like peptide-1, peptide YY, ghrelin, adiponectin and tumour necrosis factor-alpha. *Br J Nutr*. 2010;103:460-466.
- Lemoine M, Younes A. Histone deacetylase inhibitors in the treatment of lymphoma. *Discov Med*. 2010;10:462-470.
- Gao Z, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, Cefalu WT, Ye J. Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes*. 2009;58:1509-1517.
- Jiang W, Guo Q, Wu J, Guo B, Wang Y, Zhao S, Lou H, Yu X, Mei X, Wu C, Qiao S, Wu Y. Dual effects of sodium butyrate on hepatocellular carcinoma cells. *Mol Biol Rep*. 2012;39:6235-6242.
- Rodrigues MF, Carvalho É, Pezzuto P, Rumjanek FD, Amoêdo ND. Reciprocal modulation of histone deacetylase inhibitors sodium butyrate and trichostatin A on the energy metabolism of breast cancer cells. *J Cell Biochem*. 2015;116:797-808.
- Candido EP, Reeves R, Davie JR. Sodium butyrate inhibits histone deacetylation in cultured cells. *Cell*. 1978;14:105-113.
- Ansorge S, Bank U, Heimburg A, Helmuth M, Koch G, Tadjé J, Lendeckel U, Wolke C, Neubert K, Faust J, Fuchs P, Reinhold D, Thielitz A, Träger M. Recent insights into the role of dipeptidyl aminopeptidase IV (DPIV) and aminopeptidase N (APN) families in immune functions. *Clin Chem Lab Med*. 2009;47:253-261.
- Mentlein R. Dipeptidyl-peptidase IV (CD26)-role in the inactivation of regulatory peptides. *Regul Pept*. 1999;85:9-24.
- Lambeir AM, Durinx C, Scharpé S, De Meester I. Dipeptidyl-peptidase IV from bench to bedside: an update on structural properties, functions, and clinical aspects of the enzyme DPP IV. *Crit Rev Clin Lab Sci*. 2003;40:209-294.
- Dang DT, Chun SY, Burkitt K, Abe M, Chen S, Havre P, Mabeesh NJ, Heath EI, Vogelzang NJ, Cruz-Correa M, Blayney DW, Ensminger WD, St Croix B, Dang NH, Dang LH. Hypoxia-inducible factor-1 target genes as indicators of tumor vessel response to vascular endothelial growth factor inhibition. *Cancer Res*. 2008;68:1872-1880.
- Bauvois B, Djavaheri-Mergny M, Rouillard D, Dumont J, Wietzerbin J. Regulation of CD26/DPP4 gene expression by interferons and retinoic acid in tumor B cells. *Oncogene*. 2000;19:265-272.
- Cordero OJ, Salgado FJ, Viñuela JE, Nogueira M. Interleukin-12 enhances CD26 expression and dipeptidyl peptidase IV function on human activated lymphocytes. *Immunobiology*. 1997;197:522-533.
- Eltzschig HK, Faigle M, Knapp S, Karhausen J, Ibla J, Rosenberger

- P, Odegard KC, Laussen PC, Thompson LF, Colgan SP. Endothelial catabolism of extracellular adenosine during hypoxia: the role of surface adenosine deaminase and CD26. *Blood*. 2006;108:1602-1610.
22. Ishibashi Y, Matsui T, Maeda S, Higashimoto Y, Yamagishi S. Advanced glycation end products evoke endothelial cell damage by stimulating soluble dipeptidyl peptidase-4 production and its interaction with mannose 6-phosphate/insulin-like growth factor II receptor. *Cardiovasc Diabetol*. 2013;12:125.
  23. Riemann D, Kehlen A, Langner J. Stimulation of the expression and the enzyme activity of aminopeptidase N/CD13 and dipeptidyl-peptidase IV/CD26 on human renal cell carcinoma cells and renal tubular epithelial cells by T cell-derived cytokines, such as IL-4 and IL-13. *Clin Exp Immunol*. 1995;100:277-283.
  24. Stefanovic V, Ardaillou N, Vlahovic P, Placier S, Ronco P, Ardaillou R. Interferon-gamma induces dipeptidylpeptidase IV expression in human glomerular epithelial cells. *Immunology*. 1993;80:465-470.
  25. Tahara N, Yamagishi S, Takeuchi M, Tahara A, Kaifu K, Ueda S, Okuda S, Imaizumi T. Serum levels of advanced glycation end products (AGEs) are independently correlated with circulating levels of dipeptidyl peptidase-4 (DPP-4) in humans. *Clin Biochem*. 2013;46:300-303.
  26. Yamabe T, Takakura K, Sugie K, Kitaoka Y, Takeda S, Okubo Y, Teshigawara K, Yodoi J, Hori T. Induction of the 2B9 antigen/dipeptidyl peptidase IV/CD26 on human natural killer cells by IL-2, IL-12 or IL-15. *Immunology*. 1997;91:151-158.
  27. Boffa LC, Gruss RJ, Allfrey VG. Manifold effects of sodium butyrate on nuclear function. Selective and reversible inhibition of phosphorylation of histones H1 and H2A and impaired methylation of lysine and arginine residues in nuclear protein fractions. *J Biol Chem*. 1981;256:9612-9621.
  28. Byrne CS, Chambers ES, Morrison DJ, Frost G. The role of short chain fatty acids in appetite regulation and energy homeostasis. *Int J Obes (Lond)*. 2015;39:1331-1338.
  29. Dokmanovic M, Clarke C, Marks PA. Histone deacetylase inhibitors: overview and perspectives. *Mol Cancer Res*. 2007;5:981-989.
  30. Layden BT, Angueira AR, Brodsky M, Durai V, Lowe WL Jr. Short chain fatty acids and their receptors: new metabolic targets. *Transl Res*. 2013;161:131-140.
  31. Ma X, Ezzeldin HH, Diasio RB. Histone deacetylase inhibitors: current status and overview of recent clinical trials. *Drugs*. 2009;69:1911-1934.
  32. Tazoe H, Otomo Y, Kaji I, Tanaka R, Karaki SI, Kuwahara A. Roles of short-chain fatty acids receptors, GPR41 and GPR43 on colonic functions. *J Physiol Pharmacol*. 2008;59 Suppl 2:251-262.
  33. Vidali G, Boffa LC, Bradbury EM, Allfrey VG. Butyrate suppression of histone deacetylation leads to accumulation of multiacetylated forms of histones H3 and H4 and increased DNase I sensitivity of the associated DNA sequences. *Proc Natl Acad Sci U S A*. 1978;75:2239-2243.
  34. Vinolo MA, Rodrigues HG, Nachbar RT, Curi R. Regulation of inflammation by short chain fatty acids. *Nutrients*. 2011;3:858-876.
  35. Malin SK, Huang H, Mulya A, Kashyap SR, Kirwan JP. Lower dipeptidyl peptidase-4 following exercise training plus weight loss is related to increased insulin sensitivity in adults with metabolic syndrome. *Peptides*. 2013;47:142-147.
  36. Davie JR. Inhibition of histone deacetylase activity by butyrate. *J Nutr*. 2003;133(7 Suppl):2485S-2493S.
  37. Roy CC, Kien CL, Bouthillier L, Levy E. Short-chain fatty acids: ready for prime time? *Nutr Clin Pract*. 2006;21:351-366.
  38. den Besten G, Gerding A, van Dijk TH, Ciapaite J, Bleeker A, van Eunen K, Havinga R, Groen AK, Reijngoud DJ, Bakker BM. Protection against the metabolic syndrome by guar gum-derived short-chain fatty acids depends on peroxisome proliferator-activated receptor  $\gamma$  and glucagon-like peptide-1. *PLoS One*. 2015;10:e0136364.
  39. Tolhurst G, Heffron H, Lam YS, Parker HE, Habib AM, Diakogiannaki E, Cameron J, Grosse J, Reimann F, Gribble FM. Short-chain fatty acids stimulate glucagon-like peptide-1 secretion via the G-protein-coupled receptor FFAR2. *Diabetes*. 2012;61:364-371.
  40. Augustyns K, Bal G, Thonus G, Belyaev A, Zhang XM, Bollaert W, Lambeir AM, Durinx C, Goossens F, Haemers A. The unique properties of dipeptidyl-peptidase IV (DPP IV/CD26) and the therapeutic potential of DPP IV inhibitors. *Curr Med Chem*. 1999;6:311-327.
  41. Viollet B, Guigas B, Leclerc J, Hébrard S, Lantier L, Mounier R, Andreelli F, Foretz M. AMP-activated protein kinase in the regulation of hepatic energy metabolism: from physiology to therapeutic perspectives. *Acta Physiol (Oxf)*. 2009;196:81-98.
  42. Hardie DG. Sensing of energy and nutrients by AMP-activated protein kinase. *Am J Clin Nutr*. 2011;93:891S-896S.
  43. Kang W, Hong HJ, Guan J, Kim DG, Yang EJ, Koh G, Park D, Han CH, Lee YJ, Lee DH. Resveratrol improves insulin signaling in a tissue-specific manner under insulin-resistant conditions only: in vitro and in vivo experiments in rodents. *Metabolism*. 2012;61:424-433.
  44. Hardie DG. AMP-activated/SNF1 protein kinases: conserved guardians of cellular energy. *Nat Rev Mol Cell Biol*. 2007;8:774-785.
  45. Chau MD, Gao J, Yang Q, Wu Z, Gromada J. Fibroblast growth factor 21 regulates energy metabolism by activating the AMPK-SIRT1-PGC-1 $\alpha$  pathway. *Proc Natl Acad Sci U S A*. 2010;107:12553-12558.
  46. Jung GS, Jeon JH, Choe MS, Kim SW, Lee IK, Kim MK, Park KG. Renoprotective effect of gemigliptin, a dipeptidyl peptidase-4 inhibitor, in streptozotocin-induced type 1 diabetic mice. *Diabetes Metab J*. 2016;40:211-221.